


SUPPLEMENTARY MATERIAL TO  
**GC–MS-based metabolomics for the detection of adulteration in  
oregano samples**


STEFAN IVANOVIĆ<sup>1</sup>, MANUELA MANDRONE<sup>2</sup>, KATARINA SIMIĆ<sup>1</sup>, MIRJANA  
RISTIĆ<sup>3</sup>, MARINA TODOSIJEVIĆ<sup>3</sup>, BORIS MANDIĆ<sup>3</sup> and DEJAN GOĐEVAC<sup>1\*</sup>


<sup>1</sup>University of Belgrade – Institute of Chemistry, Technology and Metallurgy, Department of  
Chemistry, Njegoševa 12, 11000 Belgrade, Serbia, <sup>2</sup>University of Bologna – Department of  
Pharmacy and Biotechnology, Via Irnerio, 42, 40126 Bologna, Italy and <sup>3</sup>University of  
Belgrade – Faculty of Chemistry, Studentski trg 12–16, 11000 Belgrade, Serbia

*J. Serb. Chem. Soc.* 86 (12) (2021) 1195–1203



\* Corresponding author. E-mail: dejan.godjevac@ihtm.bg.ac.rs








Fig. S-1. GC-FID chromatograms of derivatized samples. TIC – total-ion current.