

SUPPLEMENTARY MATERIAL TO
**Thermoanalytical and spectroscopic studies on medicated jellies
with perphenazine**

MIHAELA M. BUDIUL^{1,2}, MĂDĂLINA MATEESCU^{1,2*}, GABRIELA VLASE^{1,2},
TITUS VLASE^{1,2}, SIMONA BOCĂNICI¹ and IONELA A. BRADU¹

¹Research Centre for Thermal Analysis in Environmental Problems, West University of Timișoara, Pestalozzi Street 16, 300115 Timișoara, Romania and ²West University of Timisoara – ICAM – Advanced Environmental Research Institute, Timișoara, Romania

J. Serb. Chem. Soc. 89 (4) (2024) 521–537

PREPARATION OF MEDICATED JELLIES

TABLE S-I. Components of the medicated jellies calculated on the basis [%] w/w

Jellies	Alg [%]	Biopolymers	Gel [%]	Pec [%]	BenzAc [%]	CitAc [%]	NaCit [%]	Xyl [%]	Sorb [%]	Water [%]
Alg j	50	-	-	-	0.02	0.5	0.5	10	2	36.98
Gel j	-	50	-	-	0.02	0.5	0.5	10	2	36.98
Pec j	-	-	50	-	0.02	0.5	0.5	10	2	36.98




Fig. S-1. FTIR spectra of Per (SA)/Per (D).

*Corresponding author. E-mail: madalina.mateescu@e-uvt.ro