

Synthetic study on the angular triquinanes

MIHAILO N. STJEPANOVIĆ¹, ALEKSANDAR V. JANKOVIĆ², BOJAN Z. VULOVIĆ^{1#},
RADOMIR V. MATOVIĆ² and RADOMIR N. SAIČIĆ^{1,3*}

¹University of Belgrade – Faculty of Chemistry, Studentski trg 16, 11158 Belgrade, Serbia,
²ICTM – Center for Chemistry, Njegoseva 12, 11158 Belgrade, Serbia and ³Serbian Academy
of Sciences and Arts, Kneza Mihaila 35, 11000 Belgrade, Serbia

(Received 28 June, revised 1 July, accepted 24 July 2023)

Abstract: The synthesis of an angular triquinane, which could serve as a suitable platform for the synthesis of several natural products (panaginsene, silphinene, senoxydene) is described. The synthesis is based on two consecutive cyclopentene annulations, where alkenes were used as latent carbonyl functionalities (*via* Wacker reaction), and cyclopentenone annulation was effected by aldol condensation.

Keywords: cyclopentane; natural products; organic synthesis; Wacker oxidation; aldol condensation.

INTRODUCTION

Among the thousands of structurally diverse, naturally occurring sesquiterpene compounds, the subgroup of triquinanes, although not numerous, occupies a prominent place, due to peculiar structures and a range of biological activities. These polycyclic compounds are assembled from condensed cyclopentane rings and, depending on the fusion pattern, can be classified as linear, angular or propellane (Fig. 1). Triquinane structural subunits are also embedded into tetraquinane skeletons. Not surprisingly, they have attracted considerable interest from the synthetic community.¹




Fig. 1. Three types of triquinane fusion.

* Corresponding author. E-mail: rsaicic@chem.bg.ac.rs

Serbian Chemical Society member.

<https://doi.org/10.2298/JSC230627046S>

The naturally occurring angular triquinanes comprise four skeletal types, namely isocomanes, silphinanes, pentalenanes and silphiperfolanes (Fig. 2). Some of them exhibit interesting biological activities, such as subergoric acid (**1**, member of the isocomane family), which is cardiotoxic,² but also shows antiholinesterase activity,³ and hence the ability to act as antidote against Soman in mice.⁴ Oxygenated congeners of pentalenene **2**, such as pentalenolactone, show antibiotic activity.⁵ Silphinene **3** is a biosynthetic precursor of oxygenated products such as aspergilanes,⁶ or penifulvins; penifulvin A possesses significant insecticidal activity.⁷




Fig. 2. Examples of naturally occurring angular triquinanes.


We became interested in silphinene **3** and its congeners – panaginsene **4**⁸ and senoxydene **5**. These compounds share the common angular triquinane skeleton, with some differences in the position of the alkene bond and one methyl substituent. One stereochemical difference is also of note: the methyl group at C-11 may be of either *trans*-configuration (in panaginsene and senoxydene), or *cis*-oriented (in silphinene), with respect to the cyclopentene ring. Interestingly, panaginsene was initially assigned as *cis*-configured isomer **6**, which was later revised (corrected structure: **4**).⁹ Nine total syntheses of silphinene have been reported, the first one by Ito and coworkers, who used bicyclopentadiene as the starting material.¹⁰ Paquette and Leone-Bay constructed the triquinane skeleton by iterative application of 3-oxo-organocuprate 1,4-addition/aldol condensation reaction tandem.¹¹ Sternbach and coworkers relied on the intramolecular Diels–Alder reaction, followed by oxidative fragmentation of the transient tricycle to the diquinane intermediate.¹² The synthesis by Wender and Ternansky hinges on arene-olefin meta-photocycloaddition, followed by reductive cleavage of cyclopropane ring in the tetracyclic intermediate.¹³ Crimmins and Mascarella used a

tactical combination of reactions: [2+2] alkene photocycloaddition/cyclobutane fragmentation to access racemic **3**.¹⁴ Synthetic approach by Nagarajan and Rao is based on two cyclopentane annulations, the first by intramolecular Horner-Wadsworth-Emmons reaction and the second by a radical 5-*exo*-cyclization.¹⁵ Yamamura and coworkers created a tricyclic intermediate by an intramolecular cycloaddition of electrochemically generated phenoxy cation; this intermediate was then elaborated into a triquinane framework.¹⁶ Franck-Neumann, Miesch and Gross developed “the cyclobutenic way” to silphinene, which comprises the cyclopropanation of bicyclo[3.2.0.]hept-6-ene cycloadduct, followed by solvolysis of a highly strained tricyclic intermediate.¹⁷ Fraser-Reid and Dickson used the chiron approach and synthesized (–)-**3** from mannose.¹⁸ Two syntheses of panaginsene have been reported: Lee and Geum used as a key step an intramolecular cycloaddition *via* biradical intermediate,⁹ whereas Chakraborty and Singh relied on McMurry coupling for the cyclopentene ring closure.¹⁹ As for senoxydene, two syntheses have been reported: Paquette and coworkers used intramolecular ene cyclization for the stereoselective formation of the key bicyclic intermediate,²⁰ whereas Ito and coworkers exploited the common intermediate from their previous synthesis of silphinene.²¹

RESULTS AND DISCUSSION

We set out to develop a method that would allow for the efficient construction of the angular triquinane **7**, which contains the skeleton of these three congeners, and where the configuration of the C-11 methyl group could be established subsequently (*i.e.*, in late stages of synthesis), possibly giving rise to both diastereoisomers.


Our retrosynthetic analysis is blueprinted in Scheme 1. Cyclopentenone disconnection in **7** by aldol condensation transformation simplifies the target to bicyclic ketone **8**, which could be obtained from **9** by tactical combination of reactions: allylation/Wacker oxidation. Bicyclic ketone **9** should be accessible by aldolization/isomerization tandem reaction; we were aware that this step would need a study on the reaction conditions/product distribution relationship. The dicarbonyl precursor **10** of enone **9** could be prepared by Wacker oxidation of



Scheme 1. Retrosynthetic analysis of angular triquinane **7**.

3-butenylcyclopentanone derivative (not shown), on its turn obtainable from known 4,4-dimethyl-2-cyclopentene-1-one (**11**).²²


The synthesis commenced with 1,4-addition of a butenyl cuprate reagent to the known 4,4-dimethylcyclopent-2-ene-1-one (**11**), which afforded the adduct **12** (Scheme 2). This transformation was reported earlier;^{20b} however, modifying the reaction conditions – activation by trimethylsilyl chloride and lithium chloride – improved the yield considerably (82 vs. 65 % in the previous publication).²³ Wacker oxidation of **12** under an oxygen atmosphere required 72 h at 50 °C, but provided dione **10** in 84 % yield. Various reaction conditions were tried to accomplish the cyclization step (*i.e.*, **10** → **9**), including potassium hydroxide in aqueous ether, potassium *tert*-butoxide in *tert*-butanol, or in THF, pyrrolidine (alone, or in the presence of acetic acid, or *p*-TsOH), but in all cases the mixtures of conjugated (**13**) and deconjugated (**9**) product were obtained. Finally, we found that the cyclization could be performed in boiling toluene, in the presence of catalytic amounts of *p*-TsOH, to provide the desired bicyclic enone **9** in 83 % yield (accompanied with 9 % of the conjugated isomer **13**). The next step was the installation of acetyl moiety, to obtain dione **8**. From several synthons available as synthetic equivalents of “acetyl cation” we chose the structurally simplest one – allyl bromide – for two reasons: a) minimization of steric hindrance and b) the allylation product **14** should be amenable also to aldehyde **15** – an intermediate whose role is explained below (in Scheme 3). The regioselective allylation of **9** was promoted by sodium hydride in boiling THF, where single regio- and diastereoisomer **14** was obtained (85 %). It is of note that the regioselectivity in the reactions of this type is highly dependent on a substrate structure; thus, in the allylation of the structurally related bicyclic system, the less substituted α -position of the ketone required protection, in order to secure allylation at the more substituted α' -position.²⁰ The Wacker oxidation of **14** pro-



Scheme 2. Synthesis of triquinane **7**.

ceeded much faster than the previous one (*i.e.*, **12** → **10**), and provided bicyclic dione **8** in 50 % yield (accompanied with 10 % of aldehyde **15**). Upon the treatment with potassium *tert*-butoxide in *tert*-butanol, **8** cyclized into triquinane **7** (51 %), thus providing the target molecule in six steps (12 % overall yield) from the starting 4,4-dimethylcyclopent-2-ene-1-one (**11**).

The scope of sesquiterpene natural products targeted by this methodology is not limited to angular triquinanes, but extends to bridged diquinanes, *i.e.*, cedrenes. Thus, regioisomeric oxidation of intermediate **14** should provide aldehyde **15**, from which the assembly of cedrene skeleton would be accomplished by an intramolecular aldol reaction (Scheme 3). In this way, a single synthetic intermediate (*i.e.*, **14**) would provide access to several members of two distinct classes of quinane natural products.



Scheme 3. Towards α - and β -cedrene.

CONCLUSION

To summarize, the efficient synthesis of the advanced intermediate **7** for the synthesis of triquinane sesquiterpenes (panaginsene, senoxydene and silphinene) is described, which relies on iterative application of the Wacker oxidation/aldol condensation tactical combination of reactions (*i.e.*, **12** → **9**; **14** → **7**). The research aiming to the conversion of this intermediate into the target natural products, as well as extension of this methodology to the synthesis of cedrenes (according to Scheme 3), is underway in our laboratories.

EXPERIMENTAL

General experimental details

All chromatographic separations were performed on silica gel 60 (0.063–0.2 mm), Merck. Standard techniques were used for the purification of reagents and solvents. NMR spectra were recorded on Varian/Agilent 400 (^1H -NMR at 400 MHz, ^{13}C -NMR at 100 MHz in deuterated chloroform, if not otherwise stated. Chemical shifts are expressed in ppm (δ) using tetramethylsilane as internal standard, coupling constants (J) are in Hz. IR spectra were

recorded on Thermo Scientific Nicolet Summit FT-IR instrument, and are expressed in cm^{-1} . Mass spectra were obtained on Orbitrap Exploris 240 spectrometer.

The physical data and NMR spectra of the synthesized compounds are given in Supplementary material to this paper.

4-(But-3-en-1-yl)-3,3-dimethylcyclopentan-1-one (12)

To the solution of $\text{CuBr}\cdot\text{SMe}_2$ (9.7 g, 47 mmol, 2.16 eq) and LiCl (2 g, 47 mmol, 2.16 eq) in THF (50 mL) a solution of Grignard reagent prepared from magnesium turnings (1.42 g, 58.44 mmol) and 4-bromo-1-butene (6 g, 44.44 mmol) was added dropwise at -78°C , and the reaction mixture was stirred for additional 30 min. A solution of ketone **11** (2.4 g, 21.8 mmol, 1 eq.) and TMSCl (7 mL, 5.98 g, 55 mmol, 2.52 eq.) in THF (51 mL) was added dropwise over 30 min. The reaction mixture was allowed to warm to room temperature overnight, quenched with acetic acid (22 mL) and stirred for 1.5 h. The mixture was transferred into a beaker and saturated solution of NaHCO_3 (160 mL) was added in portions. The mixture was filtered through Celite and extracted three times with methylene chloride. The combined organic layer was washed with brine, and dried over anhydrous MgSO_4 , filtered, concentrated at rotovap and purified by dry-flash chromatography (petroleum ether/ethyl acetate = 7/1) to afford 2.96 g (82 %) of product **12** as a pale yellow oil.

3,3-Dimethyl-4-(3-oxobutyl)cyclopentan-1-one (10)

Ketone **12** (610 mg, 3.67 mmol, 1 eq) was added to the solution of PdCl_2 (40 mg, 0.225 mmol, 0.06 eq) and CuCl (150 mg, 1.51 mmol, 0.41 eq.) in DMF (2.8 mL) and H_2O (0.5 mL, 27.76 mmol, 7.56 eq.) under an O_2 atmosphere. The reaction mixture was stirred at 50°C for 72 h, then diluted with water and extracted with Et_2O . Organic layer was washed with water and brine, dried over anhydrous MgSO_4 , filtered, concentrated at rotovap and purified by dry-flash chromatography (petroleum ether/ethyl acetate = 7/1) to afford 561 mg (84 %) of product **10** as pale yellow oil.

3,3,6-Trimethyl-3,3a,4,6a-tetrahydropentalen-1(2H)-one (9)

$\text{TsOH}\cdot 2\text{H}_2\text{O}$ (55 mg, 0.29 mmol, 0.22 eq) was added to the solution of diketone **10** (240 mg, 1.32 mmol) in toluene (20 mL). The flask was equipped with Deak-Stark apparatus and the reaction mixture was refluxed for 5 h. The mixture was transferred into a separatory funnel, diluted with DCM, washed with saturated solution of NaHCO_3 , dried over anhydrous MgSO_4 , concentrated at rotovap and purified by dry-flash chromatography (petroleum ether/ethyl acetate = 95/5) to afford 182 mg (83 %) of product **9** ($R_f = 0.32$ (petroleum ether/ethyl acetate = 95/5)) followed by 15 mg (9 %) of conjugated ketone **13** ($R_f = 0.17$ (petroleum ether/ethyl acetate = 95/5)).

6a-Allyl-3,3,6-trimethyl-3,3a,4,6a-tetrahydropentalen-1(2H)-one (14)

A solution of ketone **9** (200 mg, 1.22 mmol, 1 eq.) in THF (3 mL) was added to the suspension of NaH (73 mg (60 %), 1.83 mmol, 1.5 eq.) in THF (1 mL), the mixture was heated to reflux and stirred for 30 min when allyl bromide (0.21 mL, 2.43 mmol, 2 eq.) was added. After 4 h, more allyl bromide was added (0.21 mL, 2.43 mmol, 2 eq.) and refluxing was continued overnight. Reaction was quenched with water, acidified with few drops of 1.5 M HCl, extracted with DCM, washed with brine, dried over anhydrous MgSO_4 , filtered, concentrated at rotovap and purified by dry-flash chromatography (petroleum ether/ethyl acetate = 95/5) to afford 213 mg (85 %) of product **14** as colourless oil.

3,3,6-Trimethyl-6a-(2-oxopropyl)-3,3a,4,6a-tetrahydropentalen-1(2H)-one (8)

Ketone **14** (64 mg, 0.31 mmol, 1 eq.) was added to the solution of PdCl₂ (16.6 mg, 0.09 mmol, 0.3 eq.) and CuCl (40.3 mg, 0.41 mmol, 1.3 eq.) in DMF (0.7 mL) and H₂O (0.128 mL) under an O₂ atmosphere. The reaction mixture was stirred at 50 °C for 17 h, then diluted with water and extracted with Et₂O. Organic extract was washed with water and brine, dried over anh. MgSO₄, filtered, concentrated at rotovap and purified by column chromatography (hexane/ethyl acetate = 9/1) to afford 35 mg (51 %) of product **8** as colourless oil.

5,5,8-Trimethyl-4,5,5a,6-tetrahydrocyclopenta[c]pentalen-2(1H)-one (7)

The solution of diketone **8** (18 mg, 0.082 mmol, 1 eq.), *t*-BuOK (20 mg, 0.175 mmol, 2.14 eq.) and *t*-BuOH (0.1 mL) in THF (3.5 mL) was heated to 40 °C for 1 h. After cooling to room temperature, the reaction mixture was diluted with water and few drops of 1.5 M HCl, extracted with Et₂O, washed with saturated solution of NaHCO₃ and brine, dried over anh. MgSO₄, filtered, concentrated at rotovap and purified by column chromatography (petroleum ether/ethyl acetate = 5/1) to afford 8.5 mg (51 %) of product **7** as colourless oil.

SUPPLEMENTARY MATERIAL

Additional data and information are available electronically at the pages of journal website: <https://www.shd-pub.org.rs/index.php/JSCS/article/view/12460>, or from the corresponding author on request.

Acknowledgement. This research was supported by the Ministry of Science, Technological Development and Innovation of Republic of Serbia, Contract number: 451-03-47/2023-01/200168.

ИЗВОД

СИНТЕТИЧКА СТУДИЈА АНГУЛАРНИХ ТРИКИНАНА

МИХАИЛО Н. СТЕПАНОВИЋ¹, АЛЕКСАНДАР В. ЈАНКОВИЋ², БОЈАН З. ВУЛОВИЋ¹, РАДОМИР В. МАТОВИЋ²
и РАДОМИР Н. САИЧИЋ^{1,3}

¹Универзитет у Београду – Хемички Факултет, Студентски бр 16, 11158 Београд, ²ИХТМ – Центар за Хемију, Њеишева 12, 11158 Београд и ³Српска академија наука и уметности, Кнеза Михаила 35, 11000 Београд

Представљена је синтеза ангуларног трикинана, који може послужити као кључни интермедијер за синтезу неколико природних производа (панагинсена, силфинена, сен-оксидена). Синтеза се заснива на две узастопне анулације циклопентеновог прстена, код којих су алкени искоришћени као латентне карбонилне функционалне групе (преко Wacker реакције), а анулације циклопентенона су остварене алдолним кондензацијама.

(Примљено 28. јуна, ревидирано 1. јула, прихваћено 24. јула 2023)

REFERENCES

1. For review articles on triquinane synthesis, see: a) G. Mehta, A. Srikrishna, *Chem. Rev.* **97** (1997) 671 (<https://doi.org/10.1021/cr9403650>); b) L. A. Paquette, A. M. Doherty, *Polyquinane Chemistry*, 1st ed., Springer, Heidelberg, 1987 (<https://doi.org/10.1007/978-3-642-72598-2>); c) L. A. Paquette, De Gruyter, *Top. Curr. Chem.* **119** (1984) 1 (<https://doi.org/doi:10.1515/9783112539286>); d) L. A. Paquette, *Top. Curr. Chem.* **79** (1979) 41 (<https://doi.org/10.1007/BFb0048476>); e) M. B. Chanon R; Baralotto, C; Julliard, M; Hendrickson, J B, *Synthesis (Stuttg.)* **11** (1998) 1559 (<https://doi.org/10.1055/s-1998-2191>); For a recent review article on the synthesis of

- tetraquinanes, see: f) S. Kotha, A. Fatma, *Asian J. Org. Chem.* **11** (2022)e202100595 (<https://doi.org/10.1002/ajoc.202100595>)
2. a) A. Groweiss, W. Fenical, H. Cun-heng, J. Clardy, W. Zhongde, Y. Zhongnian, L. Kanghou, *Tetrahedron Lett.* **26** (1985) 2379–2382 ([https://doi.org/10.1016/S0040-4039\(00\)94832-9](https://doi.org/10.1016/S0040-4039(00)94832-9)); b) See also: V. Dragojlovic, *Molecules* **5** (2000) 674 (<https://doi.org/10.3390/50400674>)
 3. B.-H. Chen, K.-F. Jiao, Q.-E. Ji, H.-Q. Song, *J. Mol. Struct. THEOCHEM* **188** (1989) 167 ([https://doi.org/10.1016/0166-1280\(89\)85035-3](https://doi.org/10.1016/0166-1280(89)85035-3))
 4. X. Tan, H. Ye, L. Zeng, Z. Cui, S. He, *Zhongguo Haiyang Yaowu* **9** (1990) 11 (*Chemical Abstracts* 1991, 115: 35564g) [C.A 115/1991 35564g]
 5. D. E. Cane, *Sesquiterpene Biosynthesis: Cyclization Mechanisms in Compr. Nat. Prod. Chem.*, Sir D. Barton, K. Nakanishi, O. Meth-Cohn, Eds., Pergamon Press, Oxford, 1999, pp. 155–200 (<https://doi.org/10.1016/B978-0-08-091283-7.00039-4>)
 6. a) I. R. George, M. López-Tena, A. P. Sundin, D. Strand, *Org. Lett.* **23** (2021) 3536 (<https://doi.org/10.1021/acs.orglett.1c00955>); b) T. Kobayashi, Y. Teshigahara, M. Sakakibara, K. Murakami, Y. Kawamoto, H. Ito, *Org. Lett.* **25** (2023) 4510 (<https://doi.org/10.1021/acs.orglett.3c01530>)
 7. S. H. Shim, J. B. Gloer, D. T. Wicklow, *J. Nat. Prod.* **69** (2006) 1601 (<https://doi.org/10.1021/np060327z>)
 8. R. Richter, S. Basar, A. Koch, W. A. König, *Phytochemistry* **66** (2005) 2708 (<https://doi.org/10.1016/j.phytochem.2005.09.012>)
 9. S. Geum, H.-Y. Lee, *Org. Lett.* **16** (2014) 2466 (<https://doi.org/10.1021/ol500849m>)
 10. T. Tsunoda, M. Kodama, S. Itô, *Tetrahedron Lett.* **24** (1983) 83 ([https://doi.org/10.1016/S0040-4039\(00\)81333-7](https://doi.org/10.1016/S0040-4039(00)81333-7))
 11. L. A. Paquette, A. Leone-Bay, *J. Am. Chem. Soc.* **105** (1983) 7352 (<https://doi.org/10.1021/ja00363a024>)
 12. D. D. Sternbach, J. W. Hughes, D. F. Burdi, B. A. Banks, *J. Am. Chem. Soc.* **107** (1985) 2149 (<https://doi.org/10.1021/ja00293a053>)
 13. P. A. Wender, R. J. Ternansky, *Tetrahedron Lett.* **26** (1985), 2625 ([https://doi.org/10.1016/S0040-4039\(00\)98120-6](https://doi.org/10.1016/S0040-4039(00)98120-6))
 14. a) M. T. Crimmins, S. W. Mascarella, *J. Am. Chem. Soc.* **108** (1986) 3435 (<https://doi.org/10.1021/ja00272a044>); b) M. T. Crimmins, S. W. Mascarella, *Tetrahedron Lett.* **28** (1987) 5063 ([https://doi.org/10.1016/S0040-4039\(00\)95590-4](https://doi.org/10.1016/S0040-4039(00)95590-4))
 15. Y. K. Rao, M. Nagarajan, *Tetrahedron Lett.* **29** (1988) 107 ([https://doi.org/10.1016/0040-4039\(88\)80029-7](https://doi.org/10.1016/0040-4039(88)80029-7))
 16. Y. Shizuri, M. Ohkubo, S. Yamamura, *Tetrahedron Lett.* **30** (1989) 3797 ([https://doi.org/10.1016/S0040-4039\(01\)80658-4](https://doi.org/10.1016/S0040-4039(01)80658-4))
 17. M. Franck-Neumann, M. Miesch, L. Gross, *Tetrahedron Lett.* **32** (1991) 2135 ([https://doi.org/10.1016/S0040-4039\(00\)71256-1](https://doi.org/10.1016/S0040-4039(00)71256-1))
 18. J. K. Dickson, Jr., B. Fraser-Ried, *J. Chem. Soc. Chem. Commun.* (1990) 1440 (<https://doi.org/10.1039/C39900001440>)
 19. V. K. Singh, T. K. Chakraborty, *Chem. Asian J.* **16** (2021) 753 (<https://doi.org/10.1002/asia.202100144>)
 20. a) L. A. Paquette, R. A. J. Galemme, J. P. Springer, *J. Am. Chem. Soc.* **105** (1983) 6975 (<https://doi.org/10.1021/ja00361a046>); b) L. A. Paquette, R. A. Galemme Jr, J. C. Caille, R. S. Valpey, *J. Org. Chem.* **51** (1986) 686 (<https://doi.org/10.1021/jo00355a019>)
 21. T. Tsunoda, Y. Kabasawa, S. Itô, M. Kodama, *Tetrahedron Lett.* **25** (1984) 773 ([https://doi.org/10.1016/S0040-4039\(01\)80023-X](https://doi.org/10.1016/S0040-4039(01)80023-X))

22. a) P. D. Magnus, M. S. Nobbs, *Synth. Commun.* **10** (1980) 273 (<https://doi.org/10.1080/00397918008062750>) ; b) D. Pauley, F. Anderson, T. Hudlicky, *Org. Synth.* **67** (1989) 121 (<https://doi.org/10.15227/orgsyn.067.0121>) ; c) J. Yang, Y. O. Long, L. A. Paquette, *J. Am. Chem. Soc.* **125** (2003) 1567 (<https://doi.org/10.1021/ja021177r>)
23. J. Richers, A. Pöthig, E. Herdtweck, C. Sippel, F. Hausch, K. Tiefenbacher, *Chem. Eur. J.* **23** (2017) 3178 (<https://doi.org/10.1002/chem.201605362>).