



## ACCEPTED MANUSCRIPT

This is an early electronic version of an as-received manuscript that has been accepted for publication in the Journal of the Serbian Chemical Society but has not yet been subjected to the editing process and publishing procedure applied by the JSCS Editorial Office.

Please cite this article as A. M. Milošković, M. D. Radenković, N. M. Kojadinović, T. Z. Veličković, S. R. Đuretanović, and V. M. Simić, *J. Serb. Chem. Soc.* (2024) <https://doi.org/10.2298/JSC240110044M>

This “raw” version of the manuscript is being provided to the authors and readers for their technical service. It must be stressed that the manuscript still has to be subjected to copyediting, typesetting, English grammar and syntax corrections, professional editing and authors’ review of the galley proof before it is published in its final form. Please note that during these publishing processes, many errors may emerge which could affect the final content of the manuscript and all legal disclaimers applied according to the policies of the Journal.





*J. Serb. Chem. Soc.* **00(0)** 1-14 (2024)  
JSCS-12768

**Potentially toxic elements in pikeperch (*Sander lucioperca* L.) from the Gruža reservoir: Health risk assessment related to fish consumption by the general population and fishermen**

ALEKSANDRA M. MILOŠKOVIĆ<sup>1\*</sup>, MILENA D. RADENKOVIĆ<sup>2</sup>, NATAŠA M. KOJADINOVIĆ<sup>2</sup>, TIJANA Z. VELIČKOVIĆ<sup>2</sup>, SIMONA R. ĐURETANOVIĆ<sup>2</sup>, AND VLADICA M. SIMIĆ<sup>2</sup>

<sup>1</sup>University of Kragujevac, Institute for Information Technologies, Department of Sciences, Kragujevac, Serbia, and <sup>2</sup>University of Kragujevac, Faculty of Sciences, Institute of Biology and Ecology, Kragujevac, Serbia.

(Received 10 January; revised 17 January; accepted 6 April 2024)

**Abstract:** The aim was to evaluate concentrations of 14 potentially toxic elements in three tissues (muscle, liver, and gills) of pikeperch (*Sander lucioperca*) and to assess health risk (the potential non-carcinogenic - Total target hazard quotient (TTHQ) and carcinogenic - Target carcinogenic risk factor (TR) health risk associated with the consumption of pikeperch from the Gruža Reservoir by the general population and fishermen. A value of Fulton's condition factor (CF) of less than one in our study indicated the poor general health of pikeperch. According to metal pollution index (MPI), the liver was exposed to the highest pressure of metal pollution. Levels of elements were lower than the national levels and international threshold levels, and suggested a very likely absence of risk of contamination of fish with elements in the Gruža Reservoir. Higher TTHQ was observed for fishermen (0.25) compared to the general population (0.20). Higher value of TR for As compared to TR for Pb was detected, both for the general population and for fishermen. In general, there was no risk to human health from pikeperch consumption, but fishermen were at slightly higher health risk to develop cancer if they consume pikeperch meat compared to the general population.

**Keywords:** water supply reservoir; piscivore fish; fish tissues; Serbia.

#### INTRODUCTION

Numerous health benefits from consuming fish that provide many essential nutrients such as high-value proteins, various vitamins and minerals, and polyunsaturated omega-3 fatty acids, and the danger of excessive intake of potentially toxic elements, such as arsenic, mercury, cadmium, and lead, due to the

\* Corresponding author. E-mail: [aleksandra@uni.kg.ac.rs](mailto:aleksandra@uni.kg.ac.rs)  
<https://doi.org/10.2298/JSC240110044M>

consumption of contaminated fish meat are in confrontation.<sup>1</sup> Fish and fish products are among the food categories that contribute most to human exposure to dietary contaminants.<sup>2</sup> In an era where the emphasis is on healthier principles in human nutrition based on fish consumption instead of other types of meat, it is important to assess the risk of potentially toxic elements (PTEs) of the fish used in the human diet.

The globally present problem of PTEs water pollution has not bypassed Serbia, and Teodorović<sup>3</sup> highlights a large number of „hot spots“ of extreme pollution. Among aquatic biota, fish species are the most sensitive taxa to the long-term effects of pollution.<sup>4</sup> The uptake and bioaccumulation of PTEs in fish depend on the biological characteristics of fish (e.g., length and weight, age, behaviour, or nutrition), the properties of PTEs as well as properties of aquatic ecosystems.<sup>5,6</sup> Inland waters are sinks for pollutants (urban, industrial, and agricultural runoff), and according to Brönmark and Hansson<sup>7</sup> stagnant waters (reservoirs) are usually impacted by PTEs due to point sources. Lentic ecosystems (i.e., rivers) often carry concentrations of PTEs under detection limits compared to lotic (i.e., reservoirs).<sup>8</sup> The problem of PTEs pollution is more present in reservoirs due to lower self-purification capacity and pollutant dispersion in those ecosystems.<sup>9</sup>

Gruža Reservoir is located in central Serbia. This reservoir was formed by building a dam on the Gruža River in 1984, with the main purpose of supplying drinking water to the city of Kragujevac and its surroundings. Contradictory, the reservoir is also used for recreational purposes. It represents the largest water surface in Central Serbia, with an area of 934 ha. With a low water depth (average reservoir depth of 6.5 m), more than two-thirds of the reservoir has the characteristics of a lowland reservoir.<sup>10</sup> The maximum depth of the reservoir is 31 m. The reservoir suffers a strong anthropogenic influence. It is surrounded by an agricultural area where agricultural measures in the form of pesticides and herbicides are constantly applied. The reservoir also receives unprocessed wastewater from illegally built surrounding touristic settlements.

A study on the accumulation of Fe, Pb, Cd, Cu, Mn, Hg, and As in water, sediment, five macrophytes (*Typha angustifolia*, *Iris pseudacorus*, *Polygonum amphibium*, *Myriophyllum spicatum*, and *Lemna gibba*), and muscle tissue of five fish species (*Sander lucioperca*, *Abramis brama*, *Carassius gibelio*, *Silurus glanis*, and *Arystichtys nobilis*) has already been carried out in order to investigate the level of pollution in the reservoir.<sup>11</sup> The results of this study indicated higher concentrations of all examined elements in sediment than in water. Among the examined fish species, pikeperch (*Sander lucioperca*) showed the highest tendency to accumulate Pb and Hg in muscle tissue. The lack of data on the distribution of PTEs in tissues of pikeperch as the most valuable fish species in Gruža Reservoir, as well as the potential health risks, is the reason for conducting our research.

Widely distributed in Europe and Asia pikeperch (*Sander lucioperca* L.) is an indigenous fish of the Danube basin.<sup>12</sup> As a common piscivore in fish communities of many European lakes with low water transparency,<sup>13</sup> it also inhabits almost all eutrophic lakes in Serbia. As a member of the first quality group, pikeperch is an extremely valued fish species in Serbia.<sup>12,14</sup> Additionally, pikeperch is highly desirable for human consumption due to its nutritional characteristics, including the composition of proteins and fatty acids, and low-fat content (1-2%) in muscle tissue.<sup>15</sup>

The catch of fish by recreational fishermen is 1.5 higher than commercial fishing in Serbia.<sup>16</sup> When it comes to pikeperch, there is also a decline in commercial fishing. On the other hand, this species is particularly interesting for recreational fishermen. Illegal fishing of this species is also evident in the Gruža Reservoir, due to the meat's quality and high market price. Pikeperch is an important fish species in the diet, and certainly the entire catch from the Gruža Reservoir is used for human consumption.

Having in mind all of the above, this study aimed to evaluate in more detail concentrations of Al, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Sn, and Zn in three tissues (muscle, liver, and gills) of pikeperch. Also, the main aim of this study was to assess health risks (the potential non-cancerogenic TTHQ and cancerogenic TR health risk) associated with the consumption of pikeperch from the Gruža Reservoir by the general population and fishermen.

## EXPERIMENTAL

### *Fish sampling and sample preparation*

The field study was conducted at the Gruža Reservoir in central Serbia in the autumn of 2013. The sampling site coordinates are 43.927888 N, 20.678524E (Fig. 1).




Fig. 1. Map of the sampling site (43.927888N, 20.678524E) at the Gruža Reservoir

Fish (n = 20) were sampled using standing gillnets (50 m - 30 mm mesh size, 130 m - 50 mm mesh size, 100 m - 100 mm mesh size) that were left overnight. Immediately after removing the nets from the water, each pikeperch individual was sacrificed with a quick blow to the head and then dissected. Before dissection, total length (TL; to the nearest mm) and body weight (BW; to the nearest g) were measured. The measure of fish health was assessed using Fulton's condition factor (CF) with the following formula by Ricker:<sup>17</sup>

$$CF = (BW/TL^3) \times 100 \quad (1)$$

Fish dissection was done with a decontaminated ceramic knife. Tissue samples (right dorsal muscle below the dorsal fin, right gills - second arch, and liver) were washed with distilled water and transported on ice in a portable hand-held refrigerator to the laboratory.

In the laboratory, samples were weighed using an electronic scale (accuracy  $\pm 0.01$  g) and stored at  $-20$  °C prior to analysis. Before digestion in microwave Christ Alpha 2-4 LD, Harz, Germany, samples were dried in a lyophilizer Christ Alpha 2-4 LD, Harz, Germany, and measured one more time. Dried sample portions between 0.3 and 0.5 g were digested with a mixture of 65% nitric acid and 30% hydrogen peroxide (Suprapur®, Merck, Darmstadt, Germany, 10:2, v/v) at 200 °C for 20 min. After cooling to room temperature and without filtration, the solution was diluted to a fixed volume of 25 ml with ultrapure water.

#### Potentially toxic element analysis

The concentration of elements in tissues of pikeperch was measured using an inductively-coupled plasma optical emission spectrometry (ICP-OES), using a Thermo Fisher Scientific iCAP 6500 Duo ICP (Cambridge, United Kingdom). The following wavelengths were used for the ICP-OES analysis (nm): Al 391.402, As 188.032, Cd 226.602, Co 221.618, Cr 204.542, Cu 322.764, Fe 257.921, Hg 183.940, Mn 260.353, Ni 234.606, Pb 222.354, Se 199.093, Sn 245.162, and Zn 207.194. Standard muscle reference material (DORM-4, National Research Council of Canada) was digested and analyzed in triplicate to support quality assurance and control. The following assigned/measured values for DORM-4 reference material in mg kg<sup>-1</sup> are given in Table 1. Recovery ranged from 95.6 to 107.14%.

Table 1. Certified values of reference material DORM-4 and values experimentally obtained. Data are mean  $\pm$  SD

Elements	Certified values (mg kg <sup>-1</sup> )	Results obtained (mg kg <sup>-1</sup> )	Recovery (%)
As	6.80 $\pm$ 0.64	6.62 $\pm$ 0.48	97.35
Cd	0.306 $\pm$ 0.15	0.323 $\pm$ 0.019	105.56
Cr	15.9 $\pm$ 0.9	15.20 $\pm$ 0.4	95.60
Fe	341 $\pm$ 27	333.96 $\pm$ 25.87	97.93
Pb	0.416 $\pm$ 0.053	0.443 $\pm$ 0.011	106.49
Hg	0.410 $\pm$ 0.055	0.400 $\pm$ 0.064	97.56
Ni	1.36 $\pm$ 0.22	1.35 $\pm$ 0.45	99.26
Se	3.56 $\pm$ 0.34	3.61 $\pm$ 0.59	101.40
Sn	0.056 $\pm$ 0.010	0.060 $\pm$ 0.013	107.14

Mean values and standard deviations were calculated for each group, and element concentrations were expressed as mg kg<sup>-1</sup> dry weight (dw). These concentrations were recalculated to wet weight (mg kg<sup>-1</sup> ww) element concentrations, which were used to calculate

the metal pollution index (MPI), compare the concentrations of Cd, Hg, Pb, As, Cu, and Zn in fish muscles with maximum permissible concentrations (MPC) in fish meat determined by the national legislation of Serbia<sup>18</sup> and the European Union,<sup>19</sup> and to assess the risk to human health. According to these legislations, the MPCs for As, Cd, Cu, Hg, Pb, and Zn are 2.0, 0.05, 30.0, 0.50, 0.30, and 100.0 mg kg<sup>-1</sup> ww, respectively.

#### Metal pollution index (MPI)

The MPI was calculated to compare the total metals content of fish muscles, gills, and liver with each other using the following equation by Usero *et al.*:<sup>20</sup>

$$\text{MPI} = (c_1 \times c_2 \times \dots \times c_n)^{1/n} \quad (2)$$

where *c* is the concentration of the metal *n* in the sample (mg kg<sup>-1</sup> ww).

#### Liver/Muscle Hg index - L<sub>Hg</sub>/M<sub>Hg</sub>

The liver/muscle Hg index was calculated as the ratio of the concentration of Hg in the liver and muscles.<sup>21</sup>

$$\text{LHg}/\text{MHg} = \text{CL}(\text{Hg})/\text{CM}(\text{Hg}) \quad (3)$$

#### Se:Hg molar ratio

The Se:Hg molar ratio was calculated using the method of Burger *et al.*<sup>26</sup> The molar concentration of Hg was calculated by dividing the concentration of Hg (in mg kg<sup>-1</sup> ww) from muscle tissue by the molecular weight of Hg (200.59). The molar concentration of Se was calculated by dividing the concentration of Se (in mg kg<sup>-1</sup> ww) from muscle tissue by the molecular weight of Se (78.9).

#### Health risk assessments

##### Target hazard quotient - THQ

The THQ, a methodology taken from the US EPA Region III Risk-based Concentration table,<sup>22</sup> is described by the following equation:

$$\text{THQ} = \frac{(\text{EF} \times \text{ED} \times \text{FIR} \times \text{C})}{(\text{RFD} \times \text{WAB} \times \text{TA})} \times 10^{-3} \quad (4)$$

where EF is the exposure frequency (365 days/year); ED is the exposure duration (70 years), equivalent to the average lifetime; FIR is the food ingestion rate for freshwater fish for Serbia - 20 g/person/day for the general population and 25 g/person/day for fishermen;<sup>23</sup> C is the element concentration in pikeperch (mg kg<sup>-1</sup> ww); RFD is the oral reference dose (Hg = 0.0005 mg kg<sup>-1</sup>/day, Cd = 0.001 mg kg<sup>-1</sup>/day, Pb = 0.004 mg kg<sup>-1</sup>/day, Cu = 0.04 mg kg<sup>-1</sup>/day, Zn = 0.3 mg kg<sup>-1</sup>/day, Cr = 1.5 mg kg<sup>-1</sup>/day, Mn = 0.14 mg kg<sup>-1</sup>/day, Al = 0.0004 mg kg<sup>-1</sup>/day, As = 0.0003 mg kg<sup>-1</sup>/day, Fe = 0.04 mg kg<sup>-1</sup>/day, Co = 0.0003 mg kg<sup>-1</sup>/day, Ni = 0.02 mg kg<sup>-1</sup>/day);<sup>22, 24, 25</sup> WAB is the average body weight of an adult (70 kg); and TA is the average exposure time (365 days/year x ED).

Total THQ (TTHQ) was calculated using the following formula:

$$\text{TTHQ} = \sum \text{THQ} \quad (5)$$

##### Target carcinogenic risk factor - TR

The target carcinogenic risk factor (TR) for arsenic and lead was estimated using the equation:

$$\text{TR} = \frac{\text{EF} \times \text{ED} \times \text{FIR} \times \text{C} \times \text{CSF}_0}{\text{WAB} \times \text{TA}} \times 10^{-3} \quad (6)$$

where CSF<sub>o</sub> is the oral carcinogenic slope factor (mg/kg-day) which is 1.5 for As and 0.0085 for Pb.<sup>22</sup>

#### *Statistical analysis*

All values are expressed as mean (s) ± standard deviation (SD). At the beginning of the statistical analysis normality of data was tested using the Shapiro-Wilk test. In cases when data followed a normal distribution, we tested significant differences among groups using the one-way ANOVA, followed by Tukey's HSD posthoc test. On the contrary, we used the non-parametric Kruskal-Wallis H test, followed by the Mann-Whitney U test to assess differences among investigated groups. The significance level ( $\alpha$ ) was at 5%. All analyses were carried out using the SPSS 19.0 statistical package program for Windows (SPSS Inc., Chicago, IL, USA).

### RESULTS AND DISCUSSION

The weight of the pikeperch specimens examined was  $2\ 158.00 \pm 767.13$  g, while the total length was  $58.40 \pm 7.51$  cm. The CF was  $0.87 \pm 0.16$ . CF factor, as a measure of fish health, can be considered as a response to the quality of the environment. The CF value of less than one in our study indicates the poor general health of pikeperch in the Gruža Reservoir. According to Lafamme *et al.*,<sup>27</sup> Rajotte and Coutre,<sup>28</sup> and Zhelev *et al.*<sup>29</sup> CF decline was determined at highly contaminated sites. On the other hand, Kroon *et al.*<sup>30</sup> pointed out that CF as a biomarker should be examined in terms of its specificity and suitability. Overall, the value of the CF factor in this study can be seen as the first warning alarm of poor environmental conditions in the Gruža Reservoir. However, we cannot single out PTEs as the main and only reason for this condition.

The highest concentration of Hg was observed in muscle, the highest concentrations of As, Cd, Co, Cu, Fe, Se, and Zn in the liver, while the highest concentrations of Al, Cr, Mn, Ni, Pb, and Sn were detected in the gills (Table II). On the contrary, the lowest concentrations of Cd, Co, Cr, Cu, Fe, Mn, Se, Sn, and Zn in the muscle, the lowest concentrations of Al, Ni, and Pb in the liver, as well as the lowest concentrations of As and Hg in the gills were detected. Statistical tests showed no significant differences between pikeperch tissues in terms of Pb and Cr concentrations. A significant difference was recorded between all pikeperch tissues regarding Co, Cu, Fe, Ni, Se, and Zn concentrations (Table II). Muscle tissue contained significantly higher concentrations of Hg and significantly lower concentrations of Mn and Se compared to the other two tissues. Gills contained significantly higher concentrations of Al, and liver had significantly higher concentrations of As and Cd compared to the other two tissues.

Pikeperch muscle was the tissue with the lowest potential for PTEs bioaccumulation, which is confirmed by all, to date, performed studies on pikeperch fish species in Serbia.<sup>15,31-35</sup> According to Meena *et al.*,<sup>36</sup> the reason may be a low level of binding proteins in the muscle tissue. On the other hand, Hg has a high potential for bioaccumulation and biomagnification in food chains.<sup>37,38</sup> Predator fish species show important accumulation and indicator potential for Hg,



with the highest concentrations in muscle tissue.<sup>39,40</sup> As a top predator in the Gruža Reservoir, pikeperch accumulated Hg in significantly higher concentrations in muscle tissue compared to the other two tissues.

Se:Hg molar ratio in pikeperch from Gruža Reservoir was the highest in the gills (4 227.68), followed by the liver (353.85) and the lowest in the muscle tissue (10.35). Additionally, Se:Hg ratio in all three tissues was much higher than 1. This indicated that pikeperchs from Gruža Reservoir were protected against Hg toxicity, since Se:Hg molar ratio that is above 1 protects against toxicity of this element.<sup>41,42</sup>

The highest concentrations of Cu, Fe, and Zn were found in the liver of pikeperch, which agrees with results from studies by Mazej *et al.*<sup>43</sup> and Kenšová *et al.*<sup>44</sup> On the other hand, lower concentration of Cu, Fe, Zn, and Mn in muscle tissue compared to the other two tissues is in accordance with the findings of Subotić *et al.*<sup>45</sup> who stated that low level of these elements reflects the low level of binding proteins in this tissue.

Higher concentrations of Cd in pikeperch liver can be explained by the fact that this element has a very long elimination half-time, and therefore accumulates in large amounts in parenchymatous tissues such as liver.<sup>46</sup> Our results are not in accordance with the findings of Altındağ and Yiğit<sup>47</sup> and Mazej *et al.*,<sup>43</sup> who found no difference between Cd levels in the gills and liver. Given that the Gruža Reservoir is surrounded by agricultural land, our results are in agreement with the observation of Arumugam *et al.*<sup>48</sup> regarding the anthropogenic origin of Cd and As from agricultural fields, which dissolved in the water column remains for a long time in the environment.

According to Zhou *et al.*<sup>49</sup> and Ruelas-Inzunza *et al.*<sup>50</sup> gills are the organ with the highest tendency to accumulate Pb, which was also confirmed in our study. However the difference between the three tissues was not significant. The presence of Pb in the tissues of *S. lucioperca* is probably due to the traffic on the main road on the bridge that crosses the reservoir and the presence of motor boats on the surface of the reservoir.

Concentrations of Hg and As in the muscle tissue of pikeperch in this study were higher than in previous study<sup>11</sup> and this can be explained by the fish size because the fish specimens in this study are much larger.<sup>51</sup>

In comparison with the national legislation of Serbia<sup>18</sup> and the legislation of the European Union,<sup>19</sup> concentrations of As, Cd, Cu, Hg, Pb, and Zn in muscles of all pikeperch individuals were below the prescribed MPCs. The fact that the levels of elements As, Cd, Cu, Hg, Pb, and Zn were lower than the national and international threshold levels suggest a very likely absence of risk of contamination of fish with elements in the Gruža Reservoir. In the reservoirs Zlatar,<sup>35</sup> Bovan,<sup>32,52</sup> and Garaši<sup>15</sup> concentrations of elements in the muscle of the pikeperch were also below the MPCs. A recent study of Nikolić *et al.*<sup>53</sup> reported concentrations of Hg

and Cd above the MPC in the muscle of pikeperch in some 4<sup>+</sup> age group and emphasized biomagnification of these elements.

Table II. Elemental concentrations (mg kg<sup>-1</sup> dw) and metal pollution index (MPI) in muscle, liver, and gills of pikeperch (*Sander lucioperca*) in Gruža Reservoir. Values are presented as mean ± SD

Element	Muscle	Liver	Gills
Al	1.917 ± 0.607 <sup>a</sup>	1.098 ± 0.577 <sup>a</sup>	47.068 ± 26.512 <sup>b</sup>
As	0.766 ± 0.149 <sup>a</sup>	1.798 ± 0.302 <sup>b</sup>	0.630 ± 0.195 <sup>a</sup>
Cd	0.016 ± 0.004 <sup>a</sup>	0.312 ± 0.126 <sup>b</sup>	0.019 ± 0.009 <sup>a</sup>
Co	0.005 ± 0.002 <sup>a</sup>	0.431 ± 0.128 <sup>c</sup>	0.091 ± 0.035 <sup>b</sup>
Cr	0.719 ± 0.318	1.204 ± 0.435	1.283 ± 0.762
Cu	0.438 ± 0.080 <sup>a</sup>	7.947 ± 0.802 <sup>c</sup>	2.465 ± 0.711 <sup>b</sup>
Fe	6.391 ± 3.863 <sup>a</sup>	436.389 ± 212.493 <sup>c</sup>	222.349 ± 80.996 <sup>b</sup>
Hg	0.280 ± 0.090 <sup>b</sup>	0.026 ± 0.021 <sup>a</sup>	0.001 ± 0.001 <sup>a</sup>
Mn	0.359 ± 0.134 <sup>a</sup>	4.613 ± 0.927 <sup>b</sup>	4.657 ± 2.287 <sup>b</sup>
Ni	0.084 ± 0.080 <sup>b</sup>	0.015 ± 0.013 <sup>a</sup>	0.330 ± 0.131 <sup>c</sup>
Pb	0.870 ± 0.438	0.756 ± 0.312	1.005 ± 0.482
Se	1.163 ± 0.191 <sup>a</sup>	3.66 ± 0.494 <sup>c</sup>	1.663 ± 0.383 <sup>b</sup>
Sn	0.020 ± 0.007 <sup>a</sup>	0.697 ± 0.059 <sup>b</sup>	1.226 ± 0.531 <sup>b</sup>
Zn	17.713 ± 3.094 <sup>a</sup>	78.866 ± 9.414 <sup>c</sup>	41.464 ± 9.491 <sup>b</sup>
MPI	0.30	1.31	0.69

<sup>a,b,c</sup> different letters in row denote significant differences in element concentrations among the pikeperch tissues,  $p < 0.05$

According to MPI, the liver was exposed to the highest pressure of metal pollution (Table II). The lowest MPI value was recorded for muscle tissue. According to MPI, the gills were exposed to the higher pressure of metal pollution than muscle tissue, probably due to direct contact of gills with pollutants in the water.<sup>54</sup> This was also recorded for the same species in the Garaši Reservoir.<sup>15,53</sup> The liver of pikeperch was exposed to the highest pressure of metal pollution (highest MPI) as seen in pikeperch samples from the Zlatar Reservoir.<sup>35</sup> MPI values recorded in this study for all three pikeperch tissues were lower than in the same tissues of pikeperch from the Garaši Reservoir,<sup>15,53</sup> but higher than in the tissues of pikeperch from the Zlatar Reservoir.<sup>35</sup>

Liver/Muscle Hg index was 0.093. According to Havelková *et al.*,<sup>21</sup> in fish from heavily contaminated localities, the target organ for Hg accumulation is liver, while in fish from slightly contaminated localities, the main target organ for Hg accumulation is muscle. Consequently, a higher liver/muscle Hg index value is high in heavily contaminated sites. In our study, liver/muscle index value was low, indicating a slightly contaminated site. According to the above, we can conclude that Gruža is still a slightly polluted reservoir with Hg. Compared with pikeperch from the other researched reservoirs in Serbia, the concentration of Hg in the muscle tissue of pikeperch from Gruža Reservoir was higher than in the muscle

tissue of pikeperch from Bovan Reservoir,<sup>32</sup> but lower than in the muscle of pikeperch from Garaši Reservoir<sup>15</sup> and Zlatar Reservoir.<sup>35</sup>

Higher TTHQ was observed for fishermen (0.25) compared to the general population (0.20) (Fig. 2). Arsenic had the highest contribution to the overall TTHQ value, both in the general population and fishermen. The contribution of As to the overall TTHQ value was 71.40%. According to the results of THQ for all elements as well as TTHQ, the general population is under lower health risk compared to the fishermen.




Fig. 2. Total elemental THQ values due to consumption of pikeperch for general population and fishermen

The TTHQ values ranged from 0.20 to 0.25, which is much lower than the threshold value ( $TTHQ < 1$ ), indicating the absence of significant noncarcinogenic risk.<sup>55</sup> The values of TTHQ in our study were lower than in the study of the Garaši Reservoir, with values above 0.5<sup>15</sup> and the Zlatar Reservoir with values of 0.297 and 0.405.<sup>35</sup> In the two mentioned studies, the major contributor to TTHQ was Hg. According to the authors, the reason may be the lower reference dose for this element compared to other elements. In our study, the main contributor to TTHQ was As. Since agricultural activities are regularly carried out near the reservoir, we can assume that As originates from the uncontrolled use of pesticides and herbicides.

Higher values of TR for As compared to TR for Pb were detected, both for the general population and for fishermen (Table III). Fishermen are more susceptible to developing cancer if they consume pikeperch meat compared to the general population.

Table III. Target carcinogenic risk factor (TR) of As and Pb for the general population and fishermen due to consumption of pikeperch (*Sander lucioperca*)

	As	Pb
general population	$1.97 \times 10^{-6}$	$4.20 \times 10^{-7}$
fishermen	$2.46 \times 10^{-6}$	$5.25 \times 10^{-7}$

No cancerogenic risk due to intake of As and Pb from the meat of pikeperch from the Gruža Reservoir was recorded since the TRs for these elements were lower from  $10^{-6}$  or were equal to  $10^{-6}$ .<sup>56</sup> Compared to our results, a lower risk of cancer development due to As and Pb intake from pikeperch meat was recorded in the Garaši Reservoir.<sup>15</sup> Also, a lower risk of developing cancer due to intake of As from pikeperch meat was recorded in the Zlatar Reservoir.<sup>35</sup>

#### CONCLUSION

Based on the obtained results, we can conclude that despite obvious anthropogenic pressure in Gruža Reservoir and elevated concentrations of As and Hg in water,<sup>11</sup> pikeperch did not show contamination with PTEs. CF value indicated the poor general health of pikeperches indicating poor water quality. However, none of the elements exceeded MPCs and there was no noncancerogenic and cancerogenic risk to humans' health. Meat of pikeperch could be safely used by the general population and fishermen. Still, fishermen were at slightly higher health risk to develop cancer if they consume pikeperch meat compared to the general population. Due to the absences of analysis of age, gender, and diet of pikeperch in this study, the conclusions of this study should be viewed with caution. Further studies including this analysis are needed.

*Acknowledgements:* The study was supported by a Grant (Agreement No. 451-03-66/2024-03/200378) funded by the Serbian Ministry of Science, Technological Development, and Innovation.

#### ИЗВОД

#### ПОТЕНЦИЈАЛНО ТОКСИЧНИ ЕЛЕМЕНТИ У СМУЂУ (*Sander lucioperca* L.) ИЗ АКУМУЛАЦИЈЕ ГРУЖА: ПРОЦЕНА ЗДРАВСТВЕНОГ РИЗИКА ОПШТЕ ПОПУЛАЦИЈЕ И РИБАРА УСЛЕД КОНЗУМАЦИЈЕ

АЛЕКСАНДРА М. МИЛОШКОВИЋ<sup>1</sup>, МИЛЕНА Д. РАДЕНКОВИЋ<sup>2</sup>, НАТАША М. КОЈАДИНОВИЋ<sup>2</sup>, ТИЈАНА З. ВЕЛИЧКОВИЋ<sup>2</sup>, СИМОНА Р. ЂУРЕТАНОВИЋ<sup>2</sup>, И ВЛАДИЦА М. СИМИЋ<sup>2</sup>

<sup>1</sup>Универзитет у Крагујевцу, Институт за информационе технологије Крагујевац, Дејарман за природно-математичке науке, Крагујевац, Србија, <sup>2</sup>Универзитет у Крагујевцу, Природно-математички факултет, Институт за биологију и екологију, Крагујевац, Србија.

Циљ ове студије био је да се одреде концентрације 14 потенцијално токсичних елемената у три ткива (мишићи, јетра и шкрге) смуђа (*Sander lucioperca*) и да се процени здравствени ризик (потенцијално неканцерогени ризик - укупни циљни ризик од опасности (енг. Total target hazard quotient - ТТНҚ) и канцерогени ризик - циљни канцерогени фактор

ризика (енг. Target carcinogenic risk factor - TR) повезан са конзумацијом смуђа из акумулације Гружа. Вредност Фултоновог кондиционог индекса мања од један у нашој студији указује на лоше опште здравствено стање смуђа. Према индексу загађења металима (енг. metal pollution index - MPI) јетра је била изложена највишем притиску загађења металима. Концентрације елемената су биле ниже од прописаних националним и међународним законодавством, указујући на непостојање ризика од контаминиране рибе из акумулације Гружа. Примећен је већи ТТНQ за рибаре (0,25) у односу на општу популацију (0,20). Утврђена је већа вредност TR за As у поређењу са TR за Pb, како за општу популацију тако и за рибаре. Генерално, није забележен ризик за здравље људи услед конзумације смуђа, али су рибари под незнатно већим здравственим ризиком да развију рак у поређењу са општом популацијом.

(Примљено 10. јануара; ревидирано 17. фебруара; прихваћено 6. априла 2024.)

#### REFERENCES

1. A. C. Bosch, B. O'Neill, G. O. Sigge, S. E. Kerwath, L. C. Hoffman, *J. Sci. Food Agr.* **96** (2015) 32 (<https://doi.org/10.1002/jsfa.7360>)
2. WHO - World Health Organisation, Brief guide to analytical methods for measuring lead in blood. Inter-Organization Programme for the Sound Management of Chemicals 2011
3. I. Teodorović I, *Environ. Sci. Pollut. Res.* **16** (2009) 123 (<https://doi.org/10.1007/s11356-009-0152-2>)
4. N. Collin, A. Maceda-Veiga, N. Flor-Arnau, J. Mora, P. Fortuño, C. Vieira, A. de Sostoa, *Ecotox. Environ. Saf.* **132** (2016) 295 (<https://doi.org/10.1016/j.ecoenv.2016.06.017>)
5. M.G.M. Alam, A. Tanaka, G Allinson, L.J.B. Laurenson, F. Stagnitti, E.T. Snow, *Ecotox. Environ. Saf.* **53** (2002) 348 ([https://doi.org/10.1016/S0147-6513\(02\)00012-X](https://doi.org/10.1016/S0147-6513(02)00012-X))
6. J. Mwamburi, *Lakes and Reservoirs: Research Management* **18** (2013) 329 (<https://doi.org/10.1111/lre.12040>)
7. C. Brönmark, L. A. Hansson, *Environ. Conserv.* **29** (2002) 290 (<https://doi.org/10.1017/S0376892902000218>)
8. E. Has-Schön, I Bogut, R. Vuković, D. Galović, A. Bogut, J Horvatić J, *Chemosphere* **135** (2015) 289 (<https://doi.org/10.1016/j.chemosphere.2015.04.015>)
9. P. K. Rai, *Int. J. Phytoremediat.* **12** (2010) 226 (<https://doi.org/10.1080/15226510903563843>)
10. B. Ranković, J. Radulović, I. Radojević, A. Ostojić, LJ. Čomić, *Ecol. Model.* **221** (2010) 1239 (<https://doi.org/10.1016/j.ecolmodel.2009.12.023>)
11. A. Milošković, S. Branković, V. Simić, S. Kovačević, M. Ćirković, D. Manojlović, *B. Environ. Contam. Tox.* **90** (2013) 563 (<https://doi.org/10.1007/s00128-013-0969-8>)
12. P. Simonović: *Ribe Srbije*, NNK International, Beograd, Biološki fakultet Univerziteta u Beogradu, Beograd, Zavod za zaštitu prirode Srbije. 2001
13. H. Dörner, S. Berg, L. Jacobsen, S. Hülsmann, M. Brojerg, A. Wagner, *Hydrobiologia* **506** (2003) 427 (<https://doi.org/10.1023/B:HYDR.0000008608.22869.99>)

14. M. Kottelat, J. Freyhof, *Handbook of European freshwater fishes*, Publications Kottelat, Cornol, Switzerland and Freyhof, Berlin, Germany. 2007
15. D. Nikolić, S. Skorić, V. Poleksić, B. Rašković, *Environ. Sci. Pollut. Res.* **28** (2021) 53700 (<https://doi.org/10.1007/s11356-021-14526-w>)
16. Statistical Office of the Republic of Serbia, Statistical Yearbook of the Republic of Serbia, 2022 (Accessed on July 20, 2023) (<https://publikacije.stat.gov.rs/G2022/Pdf/G20222055.pdf>)
17. W.E. Ricker: *Computation and interpretation of biological statistics of fish populations*, *Bulletin*, Fisheries Research Board of Canada (1975) ISBN 0662014405 (<https://publications.gc.ca/site/eng/9.581563/publication.html>)
18. Official Gazette of the Republic of Serbia (2018) Nos. 22/2018 & 90/ 2018, Pravilnik o maksimalno dozvoljenim količinama ostataka sredstava za zaštitu bilja u hrani i hrani za životinje i o hrani i hrani za životinje za koju se utvrđuju maksimalno dozvoljene količine ostataka sredstava za zaštitu bilja. Prilog 5 – Maksimalno dozvoljene količine određenih kontaminanata u hrani i hrani za životinje biljnog i životinjskog porekla [Regulation on the maximum permitted residue levels of pesticides in food and animal feed and feed and animal feed for which maximum quantities of residues of pesticides are permitted. Annex 5 - Regulation on maximum allowed amounts of certain contaminants in food and feed for animals of plant and animal origin.], [In Serbian], 2018 ([www.pravno-informacioni-sistem.rs/SlGlasnikPortal/prilozi/5.html&doctype=reg&abc=cba&eli=true&eliActId=427071&regactid=427071](http://www.pravno-informacioni-sistem.rs/SlGlasnikPortal/prilozi/5.html&doctype=reg&abc=cba&eli=true&eliActId=427071&regactid=427071))
19. EC No. 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs (Text with EEA relevance), *Official Journal of the European Union No. 1881/2006 364* (2006) 5 (Accessed 14.03.2023) (<https://eurlex.europa.eu/eli/reg/2006/1881/oj>)
20. J. Usero, E. González-Regalado, I. Gracia, *Environ. Int.* **23** (199) 291 ([https://doi.org/10.1016/S0160-4120\(97\)00030-5](https://doi.org/10.1016/S0160-4120(97)00030-5))
21. M. Havelková, L. Dušek, D. Némethová, G. Poleszczuk, Z Svobodová, *Sensors* **8** (2008) 4095 (<https://doi.org/10.3390/s8074095>)
22. US EPA, Risk-based concentration table. Philadelphia PA: United States Environmental Protection Agency, Washington DC, 2000
23. FAO (Food and Agriculture Organization), National Aquaculture Sector, 2005 (Accessed 16.03.2023) (<http://www.fao.org/fshery/countrysector/en/>)
24. US EPA, Mercury study report to Congress health effects of mercury and mercury compounds, vol. V. Washington (DC), United States Environmental Protection Agency, EPA-452/ R-97-007, 1997
25. FAO WHO, *National Research Council Recommended Dietary Allowances 10th Edition*, National Academy Press Washington, DC, 1989
26. J. Burger, C. Jeitner, M. Donio, T. Pittfield, M. Gochfeld, *Sci. Total Environ.* **443** (2013) 278 (<https://doi.org/10.1016/j.scitotenv.2012.10.040>)
27. J. S. Lafamme, Y. Couillard, G. C. Campbell, A. Hontela, *Can. J. Fish. Aquat. Sci.* **57** (2000) 1692 (<https://cdnsiencepub.com/doi/abs/10.1139/f00-118>)
28. J. W. Rajotte, P. Couture, *Can. J. Fish. Aquat. Sci.* **59** (2002) 1296 (<https://cdnsiencepub.com/doi/abs/10.1139/f02-095>)
29. Z. M. Zhelev, S. V. Tsonev, P. S. Boyadziev *Acta Zool. Bulgar.* **70** (2018) 547 (<https://www.acta-zoologica-bulgarica.eu/downloads/acta-zoologica-bulgarica/2018/70-4-547-556.pdf>)

30. F. Kroon, C. Streten, S. Harries, *PLoS One* **12** (2017) e0174762 (<https://doi.org/10.1371/journal.pone.0174762>)
31. S. Subotić, S. Spasić, Ž. Višnjić-Jeftić, A. Hegediš, A. Krpo-Četković, B. Mićković, S. Skorić, M. Lenhardt, *Ecotox. Environ. Saf.* **98** (2013) 196 (<https://doi.org/10.1016/j.ecoenv.2013.08.020>)
32. A. Milošković, B. Dojčinović, S. Simić, M. Pavlović, V. Simić, *Fresen. Environ. Bull.* **23** (2014) 1884 ([https://hdl.handle.net/21.15107/rcub\\_cer\\_1591](https://hdl.handle.net/21.15107/rcub_cer_1591))
33. A. Milošković, B. Dojčinović, S. Kovačević, N. Radojković, M. Radenković, D. Milošević, V. Simić, *Environ. Sci. Pollut. Res.* **23** (2016) 9918 (<https://doi.org/10.1007/s11356-016-6207-2>)
34. D. A. Jovanović, R. V. Marković, V. B. Teodorović, D. S. Šefer, M. P. Krstić, S. B. Radulović, J. S. Ivanović Čirić, J. M. Janjić, M. Ž. Baltić, *Environ. Sci. Pollut. Res.* **12** (2017) 11383 (<https://doi.org/10.1007/s11356-017-8783-1>)
35. D. Nikolić, S. Skorić, B. Mićković, M. Nikčević, M. Smederevac-Lalić, V. Djikanović, *Environ. Sci. Pollut. Res.* **29** (2022) 50271 (<https://doi.org/10.1007/s11356-022-19472-9>)
36. R. A. A. Meena, P. Sathishkumar, F. Ameen, A. R. M. Yusoff, F. L. Gu, *Environ. Sci. Pollut. Res.* **25** (2018) 4134 (<https://doi.org/10.1007/s11356-017-0966-2>)
37. T. W. Clarkson, *Environ. Health Persp.* **100** (1993) 31 (<https://doi.org/10.1289/ehp.9310031>)
38. M. Gochfeld, *Ecotox. Environ. Saf.* **56** (2003) 174 ([https://doi.org/10.1016/S0147-6513\(03\)00060-5](https://doi.org/10.1016/S0147-6513(03)00060-5))
39. L. Dušek, Z. Svobodová, D. Janoušková, B. Vykusová, J. Jarkovský, R. Šmíd, P. Pavliš, *Ecotox. Environ. Saf.* **6** (2005) 256 (<https://doi.org/10.1016/j.ecoenv.2004.11.007>)
40. S. Zrnčić, D. Oraić, M. Čaleta, Ž. Mihaljević, D. Zanella, N. Bilandžić, *Environ. Monit. Assess.* **185** (2013) 1189 (<https://doi.org/10.1007/s10661-012-2625-x>)
41. S. A. Peterson, N. V. C. Ralston, P. D. Whanger, J. E. Oldfield, W. D. Mosher, *Environmental Bioindicators* **4** (2009) 318 (<https://doi.org/10.1080/15555270903358428>)
42. A. Cabañero, Y. Madrid, C. Cámara, *Biol. Trace Elem. Res.* **119** (2007) 195 (<https://doi.org/10.1007/s12011-007-8007-5>)
43. Z. Mazej, S. Sayegh-Petkovšek, B. Pokorny, *Arc. Environ. Con. Tox.* **58** (2010) 998 (<https://doi.org/10.1007/s00244-009-9417-5>)
44. R. Kenšová, O. Čelechovská, J. Doubravová, Z. Svobodová, *Acta Vet. Brno* **79** (2010) 335 (<https://doi.org/10.2754/avb201079020335>)
45. S. Subotić, Ž. Višnjić-Jeftić, S. Spasić, A. Hegediš, J. Krpo-Četković, M. Lenhardt, *Environ. Sci. Pollut. Res.* **20** (2013) 5309 (<https://doi.org/10.1007/s11356-013-1522-3>)
46. R. Cornelis, J. Caruso, H. Crews, K. Heumann, *Handbook of Elemental Speciation II. Species in the environment, food, medicine and occupational health*, Wiley, Chichester England, 2005
47. A. Altındağ, S. Yiğit, *Chemosphere* **60** (2005) 552 (<https://doi.org/10.1016/j.chemosphere.2005.01.009>)
48. A. Arumugam, J. Li, P. Krishnamurthy, Z. X. Jia, Z. Leng, N. Ramasamy, D. Du, *Environ. Sci. Pollut. Res.* **270** (2020) 19955 (<https://doi.org/10.1007/s11356-020-08554-1>)

49. H. Y. Zhou, R. Y. H. Cheung, K. M. Chan, M. H. Wong, *Water Research* **32** (1998) 3331 ([https://doi.org/10.1016/S0043-1354\(98\)00115-8](https://doi.org/10.1016/S0043-1354(98)00115-8))
50. J. Ruelas-Inzunza, F. Pérez-Osuna, D. García-Flores, *Environ. Monit. Assess.* **162** (2009) 251 (<https://doi.org/10.1007/s10661-009-0793-0>)
51. E. Zubcov, N. Zubcov, A. Ene, L. Biletschi, *Environ. Sci. Pollut. Res.* **19** (2012) 2238 (<https://doi.org/10.1007/s11356-011-0728-5>)
52. A. Milošković, V. Simić, *Pol. J. Environ. Stud.* **24** (2014) 199 (<https://doi.org/10.15244/pjoes/24929>)
53. D. Nikolić, V. Poleksić, A. Tasić, M. Smederevac-Lalić, V. Djikanović, B. Rašković, *Sustainability* **15** (2023) 11321 (<https://doi.org/10.3390/su151411321>)
54. P. B. Hamilton, G. Rolshausen, T. M. Uren Webster, C. R. Tyler, *Philosophical Transactions of the Royal Society of London Series B* **372** (2017) 20160042 (<https://doi.org/10.1098/rstb.2016.0042>)
55. N. Zheng, Q. Wang, X. Zhang, D. Zheng, Z. Zhang, S. Zhang, *Sci. Total Environ.* **387** (2007) 96 (<https://doi.org/10.1016/j.scitotenv.2007.07.044>)
56. M. S. Islam, M. K. Ahmed, M. Habibullah-Al-Mamun, K. N. Islam, M. Ibrahim, S. Masunaga, *Food Additives & Contaminants: Part A* **31** (2014) 1982 (<https://doi.org/10.1080/19440049.2014.974686>).