



ACCEPTED MANUSCRIPT

This is an early electronic version of an as-received manuscript that has been accepted for publication in the Journal of the Serbian Chemical Society but has not yet been subjected to the editing process and publishing procedure applied by the JSCS Editorial Office.

Please cite this article as S. S. Konstantinović, D. Z. Troter, J. B. Zvezdanović, S. M. Petrović, S. R. Savić, B. R. Danilović, and J. Ćirić, *J. Serb. Chem. Soc.* (2025) <https://doi.org/10.2298/JSC250619084K>

This “raw” version of the manuscript is being provided to the authors and readers for their technical service. It must be stressed that the manuscript still has to be subjected to copyediting, typesetting, English grammar and syntax corrections, professional editing and authors’ review of the galley proof before it is published in its final form. Please note that during these publishing processes, many errors may emerge which could affect the final content of the manuscript and all legal disclaimers applied according to the policies of the Journal.



J. Serb. Chem. Soc. **00(0)** 1-14 (2025)
JSCS-13428

Reishi (*Ganoderma lucidum*) tincture enriched with propolis or green tea from Serbia: UHPLC-DAD-ESI-MS characterization, mineral content and antibacterial activity

SANDRA S. KONSTANTINOVIC^{1*}, DRAGAN Z. TROTTER¹, JELENA B. ZVEZDANOVIC¹, SANJA M. PETROVIC¹, SAŠA R. SAVIC¹, BOJANA R. DANILOVIC¹ AND JOVAN ĆIRIĆ²

¹University of Niš, Faculty of Technology, Bulevar oslobođenja 124, 16000 Leskovac, Serbia, and ²Toplica Academy of Vocational Studies, Department of Agricultural and Food Studies, Ćirila i Metodija 1, 18400 Prokuplje, Serbia.

(Received 19 June; revised 21 July; accepted 20 November 2025)

Abstract: The objective of the present study was to evaluate the differences in chemical composition and mineral content of pure Reishi (*Ganoderma lucidum*) tincture with those enriched with propolis or green tea. The chemical composition and mineral content were analyzed by UHPLC-DAD-ESI-MS and ICP-OES methods, respectively. Among all samples, Reishi/propolis tincture contained a significant amount of polyphenols, with the dominant contribution of flavonoids and phenolic acids. Regarding mineral content, P, K, and Na were the most abundant in all tinctures. Reishi/propolis tincture also demonstrated the best antibacterial activity, with *P. vulgaris* and *P. aeruginosa* as the most sensitive bacteria.

Keywords: antibacterial activity; green tea; ICP-OES; propolis; reishi; UHPLC-DAD-ESI-MS.

INTRODUCTION

Reishi, or Lingzhi (*Ganoderma lucidum*), is one of the most popular and highly regarded medicinal fungi, demonstrating various biological and pharmacological activities, e.g., anti-tumor, antimicrobial, antioxidative, anti-inflammatory, anti-hyperglycemic, anti-hyperlipidemic, anti-tyrosinase and antimutagenic, as well as cardio-, neuro-, hepato-, and renal-protective, etc.^{1,2} Commercial *G. lucidum* products are available as powders, dietary supplements, and tea obtained from its different parts, including mycelia, spores, and fruit bodies. The main component of mushrooms is water (90% w/w), while proteins, fats, carbohydrates, ash, minerals, and vitamins are the other 10% of the total

* Corresponding author. E-mail: sakisandra12@yahoo.com
<https://doi.org/10.2298/JSC250619084K>

content.^{2,3} The fruit body of this fungus is a rich source of bioactive components, such as polysaccharides, triterpenes, polyphenols, vitamins and different types of secondary metabolites (phenolic compounds, polyketides, terpenes, steroids, and phenols), including more than 140 different lanostane-type triterpenoids.¹⁻³ Such a versatile chemical composition explains a wide range of its pharmaceutical properties.

Combining pure Reishi tincture with other natural and biologically active extracts can improve its biological and pharmacological activities. Propolis, a resinous substance collected by bees from different plants, is another natural product considered one of the best general panaceas since ancient times.⁴ It is a complex mixture, and its chemical composition is qualitatively and quantitatively variable, depending on the bee species, geographical areas, and botanical source.^{5,6} Among natural compounds found in propolis, polyphenols are the most important, mainly phenolic acids and flavonoids.⁴⁻⁶ These compounds are responsible for its numerous activities, such as antimicrobial, antioxidant, anti-inflammatory, antiproliferative, antihypertensive, spasmolytic, hepatoprotective, immune-stimulant, *etc.*⁴⁻⁶ Propolis can be found in various cosmetics, beverages, food products, and dietary supplements. Green tea, obtained from the leaves of *Camellia sinensis* L., is a widely consumed beverage worldwide. It is rich in polyphenols (particularly catechins), while its extract contains lower amounts of other compounds, such as flavonoids and phenolic acids.⁷

Thus, the aims of the current study were to determine and compare the chemical and mineral composition of pure Reishi tincture with those enriched with propolis or green tea from Serbia, as well as to evaluate their antimicrobial activity against eight bacteria. The chemical composition and mineral content were analyzed by UHPLC-DAD-ESI-MS and ICP-OES methods, respectively. These results could promote further development of the Reishi production industry in Serbia and be useful for the food, pharmaceutical, and cosmetic industries.

EXPERIMENTAL

Material

Commercial ethanolic extracts obtained by triple extraction of pure Reishi (35%), Reishi/Propolis (35%), and Reishi/Green tea (35% v/v) were analyzed. A total of 100 ml of each tincture was evaporated by rotary evaporator at 40 °C to dryness and dried at 50 °C until reaching constant weight. The total solid matter was 4 mg·mL⁻¹ of tincture.

Chemicals and reagents

Ethanol, 96% (Centrochem, Zemun, Serbia), and quercetin, rutin, and apigenin standards (Sigma-Aldrich, St. Louis, USA) were used. Distilled water and aqueous NaCl solution (0.9%) were provided from Zorka Pharma d.o.o. (Šabac, Serbia). Dimethylsulfoxide (DMSO) (≥ 99%) was purchased from Sigma-Aldrich (Munich, Germany). Solvents used in the UHPLC experiments were of HPLC or LC/MS grade. Concentrated nitric acid (65%, Merck) and distilled water (HPLC grade, Fisher Chemical) were used for ICP-OES analysis.

UHPLC-DAD-ESI-MS analysis

The chromatographic separation was performed on a Hypersil Gold C18 column (50×2.1 mm, 1.9 μm) at 25 °C using a Dionex Ultimate 3000 UHPLC+ system equipped with a diode array (DAD) detector and LCQ Fleet Ion Trap Mass Spectrometer (Thermo Fisher Scientific, USA). A mobile phase consisted of two solvents, 0.1 %, v/v, formic acid in water and methanol, with a gradient program at 0.25 mL·min⁻¹ flow rate.⁸ The injection volume was 1 μL . Absorption UV/Vis spectra were recorded on a DAD-detector with a total spectral range between 200 and 800 nm. The mass spectrometric analysis was performed using a 3D-ion trap with electrospray ionization (ESI) in negative ion mode. The ESI-source parameters were as follows: source voltage 4.5 kV, capillary voltage -41 V, tube lens voltage -95 V, capillary temperature 350 °C, nitrogen sheath and auxiliary gas flow 32 and 8 arbitrary units. MS spectra were acquired by full range acquisition of m/z 130–1300, with a tandem mass spectrometry analysis performed by a data-dependent scan – the collision-induced dissociation (CID) of detected molecular ion peaks ($[\text{M}-\text{H}]^-$) with normalized collision energy set at 25 eV. The ESI-source parameters for positive ion mode also used for detection of pure Reishi tincture were source voltage 5.0 kV, capillary voltage 49 V, tube lens voltage 100 V, capillary temperature 350 °C, and 32 and 8 units for the auxiliary and sheath gas flow, respectively, in the full range, 130–1200, and data-dependent scan with CID set at 25 eV of detected $[\text{M}+\text{H}]^+$, or cationized $[\text{M}+\text{Na}]^+$ molecular ion peaks. Xcalibur software (version 2.1) was used for instrument control, data acquisition, and data analysis.

ICP-OES analysis

The quantitative analysis was carried out using ICP-OES (Inductively Coupled Plasma - Optical Emission Spectrometry, ARCOS FHE12, SPECTRO, Germany), following the manufacturer's guidelines. All samples from the solid (Reishi, propolis, and green tea powders) and liquid (tinctures) materials were prepared through a wet digestion procedure with concentrated nitric acid, as detailed in the literature.^{9,10} Argon 5.0 (99.999%) purity served as the carrier gas. In addition to the analytical sample, calibration standards were prepared beforehand. The calibration solution was created using Multistandard IV – a multi-elemental standard solution from Merck, which includes Ag, B, Cd, Co, Cr, Cu, Fe, Li, Mn, Na, Ni, Pb, Tl, and Zn at concentrations of 1000 ppm. Parameters of the calibration lines for the determined elements are presented in Supplementary material (Table S-I.).

Antimicrobial activity

The antimicrobial activity was evaluated using the following microorganisms: *S. aureus* (ATCC 25923), *P. aeruginosa* (ATCC 27853), *E. coli* (ATCC 25922), *K. pneumoniae* (ATCC 700603), *L. monocytogenes* (ATCC 19166), *P. vulgaris* (ATCC 8427), *B. cereus* (ATCC11778) and *B. luteus* in haus strain. Microorganisms were from the Microbiology Laboratory, Faculty of Technology, Leskovac. Media used for the growth of the microorganisms were Antibiotic agar no. 1 for microbiology (Merck, Darmstadt, Germany) and nutrient agar (Torlak, Belgrade, Serbia) for bacterial growth. The agar disc diffusion method was used for testing the antibacterial activity.¹¹ The mediums were sterilized in an autoclave at 121 °C under 110 kPa for 25 min. The suspension was prepared with overnight culture and adjusted to 0.5 McFarland standard. The prepared suspension contained 1-2·10⁸ CFU·mL⁻¹ (CFU – colony-forming unit). The inoculum of 0.1 mL of suspension was added to 10 mL of medium and poured into the Petri dishes. For screening, sterilized filter paper disks (9 mm in diameter, Schleicher & Schuell) were placed on the surface of inoculated mediums and impregnated with 50 μL of the extract sample (evaporated ethanolic extract dissolved in DMSO, concentration 100 mg·mL⁻¹). Pure

DMSO was used as a negative control. Plates were incubated for 24 h at 37 °C and results were expressed as the diameter of inhibition zones (mm). The presence of the inhibition zone (IZ) indicated the activity of the tested samples against the tested microorganisms. The following commercial standardized discs with predetermined amounts of antibiotics were used as positive controls: Ciprofloxacin (5 µg), Ipinem (10 µg), Penicillin (6 µg, 10 IU), Trimethoprim/sulfamethoxazole (1.25/23.75 µg), Clindamycin (2 µg), Ceftriaxone (30 µg), Gentamicin (10 µg), Amoxicillin + clavulanate (20/10 µg), and Cefoxitin (30 µg) (BD Sensi-Disc, GmbH, Heidelberg). All experiments were performed in triplicate.

Statistical analysis

All results were reported as mean value ± standard deviation from three measurements. Statistical comparisons were carried out by one-way ANOVA (analysis of variance) followed by Tukey's multiple comparison test by software SPSS 23.0 (IBM, USA). Differences are reflected as significant at $p < 0.05$.

RESULTS AND DISCUSSION

UHPLC-DAD-ESI-MS analysis

The typical UHPLC-DAD chromatograms recorded from DAD-signal at $\lambda = 320$ nm are shown in Fig. 1(a-c).




Fig. 1. UHPLC-DAD chromatograms of all tinctures (a-pure Reishi, b-Reishi/Green tea and c-Reishi/Propolis) at DAD signal $\lambda_{\text{det}} = 300$ nm. The corresponding peaks assignments (in the chromatograms labeled by numbers) are listed in Tables S-I, S-II and S-III, respectively.

The list of all compounds detected with their MS/UV/Vis spectral data is shown in **Tables S-II–S-IV**. For some detected compounds, the absorption and MS/MS spectra are shown in Supplementary Data (**Figs. S-1 and S-2**). Mass

spectra and fragmentation patterns, the order of elution from the chromatographic system, together with literature data,¹³⁻²⁸ allowed the identification of some detected compounds.

Basic physiologically active components in *G. lucidum* are polysaccharides, peptidoglycans, and triterpenes.¹⁻³ Among 13 registered DAD-signals in pure Reishi tincture (Fig. 1a, Table S-II), signal 10 was identified as 12-acetoxylanoderic acid F, according to $[M-H]^-$, m/z 569, while ganoderic acid G type A ($[M-H]^-$, m/z 531 in negative mode and $[M+Na]^+$, m/z 555 in positive mode) was assigned to peak 12.¹⁶ Those ganoderic acids (related to triterpenoids) are the most representative components in *G. lucidum*, responsible for different biological activities. An appropriate MS/MS-fragmentation pattern in positive mode for the same compound (Fig. S-1a) confirmed the suggested compound, with cationized adducts at m/z 555 $[M+Na]^+$ and corresponding fragment ions. The corresponding UV/Vis spectrum (Fig. S-1a) of the ganoderic acid G type A exhibits an absorption maximum at 258 nm.¹⁶ The triterpenoid often present in *G. lucidum*, lucidone A, was tentatively identified by confirming with Yan *et al.*¹⁵

The dominant class of compounds in both Reishi/propolis and Reishi/green tea tinctures are flavonoids. Rutin (Fig. 1b, Table S-III; peak No. 11), quercetin (Fig. 1c, Table S-IV; peak No. 9), and apigenin (Fig. 1c, Table S-IV; peak No. 14) were identified by their standard methanolic solutions. Hydroxybenzoic acid, caffeic acid, *p*-coumaric acid, caffeic acid derivate, and dihydroquercetin (Fig. 1c, Table S-III; peaks No. 1, 2, 3, 4, and 5, respectively) were identified by using MassBank Record.¹⁹ For qualitative analysis of all detected signals, mass spectrometry combined with collision induced dissociation of molecular ions in data-dependent mode MS/MS was used, together with a comparison with the corresponding literature (Tables S-III and S-IV).

Reishi/green tea tincture is rich in catechin, epicatechin, galocatechin, epigallocatechin, and flavan-3-ols, characteristic of green tea. Galocatechin is detected as signal No. 2 in the UHPLC-DAD chromatogram (Fig. 1b, Table S-III) and confirmed by the MS/MS-fragmentation pattern in negative mode and corresponding fragment ions (Fig. 2b). The UV/Vis spectrum of galocatechin (Fig. S-1b) exhibits an absorption maximum at 287 nm.^{18,29} Reishi/green tea tincture also contains flavonols: kaempferol 3-*O*-rutinoside, kaempferol-*O*-glycoside, quercetin-glycoside and its quercetin-3-*O*-rutinoside derivative rutin (Fig. 1b, Table S-III; peaks No. 13, 14, 12, and 11, respectively) and flavones such as apigenin-glucosides: apigenin 6 (or 8)-*C*-xyloside-8(or 6)-*C*-glucoside (known as vicanin 1, IUPAC name 5,7-dihydroxy-2-(4-hydroxyphenyl)-8-[(2S,3R,4R,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]-6-[(2S,3R,4S,5R)-3,4,5-trihydroxyoxan-2-yl]chromen-4-one) and apigenin 6-*C*-glucosyl 8-*C*-(2"-*O*-dihydroferuloyl)-glucoside (IUPAC name 5,7-dihydroxy-2-(4-hydroxyphenyl)-6-[(2S, 3R, 4R, 5S, 6R)-3, 5-trihydroxy-6-

(hydroxymethyl)oxan-2-yl]-8- $\{[2R,3R,4S,5S,6R)$ -2-[3-(4-hydroxy-3-methoxyphenyl)propanoyl]oxy-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl] $\}$ chromen-4-one) (Fig. 1b, Table S-III; peaks No. 9 and 10, respectively). MS/MS fragmentation pattern in a negative mode for vicenin 1 (Fig. S-1c) presents molecular ions at m/z 563, 443, 383, and 353. These neutral losses are characteristics of glucosyl and pentosyl residues. The fragmentation signals at m/z 383 and 353 are also typical for di-*C*-glycosylflavones. UV/Vis spectra of vicenin 1 (Fig. S-1c) exhibit two absorption maxima at 272 and 334 nm, which is in accordance with the literature.²⁰ Quercetin-glycoside (Fig. 1b, Table S-III; peak No. 12) is detected as quercetin-7-*O*-glycoside, since the fragment ion at m/z 301 is rather abundant in 7-*O*-glycoside than in 3-*O*-glycoside (Fig. S-1d). Moreover, UV/Vis absorption maxima at 271 and 349 nm are in the range for this class of flavonols.²¹

Reishi/Propolis tincture is rich in phenolic acids. Caffeic acid was detected as peak No. 2 in the UHPLC-DAD chromatogram (Fig. 1c, Table S-IV), while the corresponding UV/Vis spectrum (Fig. S-2a) contains two absorption maxima at 290 and 323 nm. An appropriate MS/MS-fragmentation pattern in negative mode (Fig. S-2a) confirmed the suggested compound with molecular ion at m/z 179 $[M-H]^-$ and corresponding fragment ion at m/z 135.^{19,23} Caffeic acid derivatives were detected by peaks No. 4, 20, 21 and 22 (Fig. 1c, Table S-IV).^{19,23-25,28} Reishi/Propolis tincture also contains quercetin and its derivatives dihydroquercetin or taxifolin, quercetin-*O*-rhamnoside quercetin-3-methyl-ether and quercetin-3-dimethyl-ether (Fig. 1c, Table S-IV; peaks No. 9, 5, 7, 11 and 19) with molecular ion peaks found at m/z 301, 303, 447, 300, and 329, respectively. The corresponding UV/Vis spectrum of taxifolin (Fig. S-2b) contains two absorption maxima at 298 and 323 nm, while an appropriate MS/MS-fragmentation pattern in negative mode (Fig. S-2b) confirmed the suggested compound.^{19,25} Genistin, a well-known isoflavone, is identified as a peak No. 8 ($[M-H]^-$, m/z 431) (Fig. 1c, Table S-IV). Corresponding UV/Vis spectrum (Fig. S-2c) contains two absorption maxima at 290 and 322 nm, typical for this compound. The product ion mass spectrum of $[M-H]^-$ ions (Fig. S-2c) shows neutral losses of the glycan residue (162 Da) and of the glycan residue +H \cdot (163 Da) by rearrangement and cleavage, respectively, which is expected since, structurally, genistin represents genistein 7-*O*- β -D-glucoside.^{19,25} Reishi/propolis tincture is also rich in dihydroflavonol pinobanksin, its 5-methyl-ether derivative and 3-*O*-derivatives (acetate, butyrate, propionate, and pentanoate), as well as methoxy-chrysin, luteolin-methyl-ether, galangin-5-methyl-ether and isorhamnetin 3-*O*-rutinoside.^{24,26,28} The tincture also contains flavanone pinocembrin (Fig. 1c, Table S-IV; peak No. 23), a natural precursor of pinobanksin. The corresponding UV/Vis spectrum (absorption maxima at 291 and 356 nm) and MS/MS-fragmentation

pattern (Fig. S-2d) confirmed its presence.²⁸ Peaks No. 12, 13, 18 and 26 were unidentified derivatives of phenolic acids and flavonoids.

Mineral content

ICP-OES analysis results are presented in Table I.

TABLE I. Concentrations of macro- and microelement in samples^a

	Solid material $c / \mu\text{g}\cdot\text{g}^{-1}$			Tincture $c / \text{mg}\cdot\text{L}^{-1}$		
	Reishi	Green Tea	Propolis	Reishi	Reishi/green tea	Reishi/propolis
Ag	6.45 ^{La}	0 ^{Kb}	0 ^{Nb}	0 ^{Kb}	0 ^{Ob}	0 ^{Mb}
B	6.05 ^{Mb}	3.54 ^{Gc}	31.77 ^{Ha}	0 ^{Ke}	3.28 ^{Ed}	0 ^{Me}
Ca	0 ^{Oc}	112.1 ^{Bb}	760.43 ^{Da}	0 ^{Kc}	0 ^{Oc}	0 ^{Mc}
Cd	0 ^{Oc}	6.8 ^{Fa}	0 ^{Nc}	0 ^{Kc}	0 ^{Oc}	0.025 ^{Lb}
Co	0 ^{Ob}	0.03 ^{Ka}	0 ^{Nb}	0 ^{Kb}	0 ^{Ob}	0 ^{Mb}
Cr	44.76 ^{Ja}	0.45 ^{J,Kb}	0 ^{Nc}	0 ^{Kc}	0 ^{Oc}	0 ^{Mc}
Cu	77.02 ^{Fa}	2.45 ^{Hc}	8.49 ^{Lb}	0.15 ^{If}	0.83 ^{Jd}	0.4 ^{Ie}
Fe	1071.78 ^{Ca}	2.2 ^{Hc}	26.43 ^{Ib}	0 ^{Ke}	0.43 ^{Ld}	0 ^{Me}
K	2.02 ^{Nf}	11 ^{Ee}	2659.07 ^{Ba}	89.78 ^{Bb}	49.33 ^{Bc}	26.25 ^{Cd}
Li	174.6 ^{Ea}	2.2 ^{Hb}	1.21 ^{Mc}	0.1 ^{Je}	0.13 ^{Md}	0.1 ^{Ke}
Mg	0 ^{Oe}	320 ^{Ab}	383.12 ^{Ea}	4.9 ^{Ec}	2.95 ^{Gd}	2.05 ^{Fd}
Mn	739.11 ^{Da}	0.5 ^{J,Kc}	15.28 ^{Jb}	0 ^{Ke}	0.075 Nd	0 ^{Me}
Na	0 ^{Of}	93 ^{Cb}	318.14 ^{Fa}	35.6 ^{Cd}	48.23 ^{Cc}	34.48 ^{Ae}
Ni	13.31 ^{Ka}	2.35 ^{Hb}	0 ^{Nc}	0 ^{Kc}	0 ^{Oc}	0 ^{Mc}
Pb	45.16 ^{Ia}	1.15 ^{Jc}	0 ^{Ne}	0 ^{Ke}	1.45 ^{Ib}	0.25 ^{Jd}
Sr	0 ^{Oe}	0 ^{Ke}	11.4 ^{Ka}	0.53 ^{Hc}	0.73 ^{Kb}	0.5 ^{Hd}
Tl	47.18 ^{Ha}	0.53 ^{J,Kb}	0 ^{Nc}	0 ^{Kc}	0 ^{Oc}	0 ^{Mc}
Zn	48.79 ^{Ga}	0.3 ^{Kf}	36.13 ^{Gb}	2.38 ^{Gc}	2.35 ^{Hd}	1.7 ^{Ge}
Si	5657.26 ^{Aa}	2.25 ^{Hf}	35.89 ^{Gb}	3.45 ^{Fd}	3.08 ^{Fe}	3.85 ^{Ec}
P	1594.76 ^{Bb}	0.88 ^{I,Jf}	8692.05 ^{Aa}	238.48 ^{Ac}	117.58 ^{Ad}	31.15 ^{Be}
S	0 ^{Of}	29.25 ^{Db}	1244.91 ^{Ca}	26.2 ^{Dc}	21.88 ^{Dd}	6.53 ^{Be}

^a column values (different capital letters) and row values (different small letters) are significantly different ($p < 0.05$) by Tukey's multiple range test

The difference in mineral content between crude samples and tinctures is evident, not only in a qualitative but also in a quantitative manner. A total of 21 elements was determined. Tested Reishi powder is notable for its high content of Si (5657.26 $\mu\text{g}\cdot\text{g}^{-1}$), P (1594.76 $\mu\text{g}\cdot\text{g}^{-1}$), Fe (1071.78 $\mu\text{g}\cdot\text{g}^{-1}$), and Mn (739.11 $\mu\text{g}\cdot\text{g}^{-1}$), while propolis was rich in P (8692.05 $\mu\text{g}\cdot\text{g}^{-1}$), K (2659.07 $\mu\text{g}\cdot\text{g}^{-1}$), S (1244.91 $\mu\text{g}\cdot\text{g}^{-1}$), and Ca (760.43 $\mu\text{g}\cdot\text{g}^{-1}$), followed by Mg (383.12 $\mu\text{g}\cdot\text{g}^{-1}$) and Na (318.14 $\mu\text{g}\cdot\text{g}^{-1}$). Among all crude samples, green tea powder contained the least amount of minerals, with Mg as the most abundant element (320 $\mu\text{g}\cdot\text{g}^{-1}$), followed by significantly lower quantities of Ca (112.1 $\mu\text{g}\cdot\text{g}^{-1}$), Na (93 $\mu\text{g}\cdot\text{g}^{-1}$), and S (29.25 $\mu\text{g}\cdot\text{g}^{-1}$). Although the mineral content of plants depends upon soil structure, climatic conditions, and plant species, the basic elements are more or less present in the same species. According to literature, Reishi contains minerals like Cu, Zn,

Se, K, Ca, P, Mg, and Fe.³⁰ Propolis possesses Ca, K, Al, Cu, Mg, Fe, and Zn in the highest amounts.³¹ The most abundant minerals found in green tea are P, K, Ca, Mg, Fe, Mn, Al, S, Si, followed by Zn, Cu, and F.³² In all Reishi tinctures, the most common minerals are P, K, and Na, which are very important macroelements and have a large impact on the human body. P plays a crucial and vital part in the creation of bones, teeth, nucleic acids, phosphoproteins, growth, and the control of different repair processes. Its role in the conversion of food into energy, in the metabolism of lipids and carbohydrates, in lessening arthritis pain, and in restoring proper kidney and cardiac function is also essential.³³ As two essential macroelements, K and Na are important to maintain body cellular homeostasis and metabolism, since most metabolic processes are dependent on or affected by these electrolytes. Also, these electrolytes maintain normal pH levels in extracellular fluid, regulate osmotic pressure, act as cofactors for many enzymes, and are involved in oxidation-reduction reactions.³⁴ Compared to crude samples, in all Reishi extracts, toxic elements were not found or were present in insignificant amounts, such as Cd and Pb.

Antimicrobial activity

The results of the disc diffusion testing are presented in Table II. The DMSO or sterile water negative controls showed no inhibition effects. The inhibition zones reveal different susceptibility of different bacteria to all tinctures.

None of the tinctures was antibacterial against *K. pneumonia*. While pure green tea tincture failed to express any antibacterial activity, both the pure Reishi and propolis tinctures demonstrated similar potential against *P. aeruginosa*, *E. coli*, and *P. vulgaris*. However, propolis tincture was more active against *S. aureus*, and was the only pure tincture deemed as antibacterial against *L. monocytogenes*, *B. cereus* and *B. luteus*. The antibacterial potential of different Reishi tinctures is reported in earlier studies.^{35,36}

Compared to the pure Reishi extract, the Reishi/green tea tincture exhibited improved antibacterial effect, while Reishi/propolis tincture showed the highest activity against all tested strains, especially against *S. aureus* and *L. monocytogenes*. Somewhat improved antibacterial activity of Reishi/green tea tincture compared to the pure Reishi and pure green tea tinctures can only be explained by the possible synergistic effect of the components present in Reishi and green tea. Specifically, the main catechins present in green tea (epicatechin, epicatechin-3-gallate, epigallocatechin and epigallocatechin-3-gallate) are known antimicrobials³⁷; these catechins are detected in prepared Reishi/green tea tincture.

Interestingly, the addition of propolis in the Reishi tincture enabled it to express activity against *L. monocytogenes*, *B. luteus* and *B. cereus*. Previous studies confirmed the antibacterial potential of propolis tinctures from Serbia against *L. monocytogenes*, *S. aureus* and *E. coli*.³⁸⁻⁴⁰ Although the chemical composition of propolis is quite complex, its source causes only minor variations

regarding its antimicrobial activity.⁴⁰ Antimicrobial activity of propolis was associated with the presence of phenolic compounds, such as caffeic acid, quercetin, luteolin, apigenin, *p*-coumaric acid, kaempferol, and pinobanksin.³⁹ Since these phenolic components are also present in Reishi/propolis tincture, its improvement in antibacterial capacity is explained by both the presence of various phenolic compounds originating from Reishi and propolis, as well as their synergistic effect.

Although the Reishi/propolis tincture exhibits moderate antibacterial activity compared to some commercial options, its use may still be justified due to the multiple other potential health benefits associated with its constituents. Previously reported immunomodulatory, anti-inflammatory, and antioxidant properties of both Reishi¹⁻³ and propolis⁴⁻⁶ could contribute to overall wellness.

TABLE II. Values of zones of bacterial growth inhibition (mm)^a

Tincture	SA	PA	EC	KP	LM	BC	BL	PV
R	9*±0.57 ^{Jc}	12±0.5 ^{Hb}	10±0.5 ^{Gc}	NI ^{Gd}	NI ^{Hd}	NI ^{Fd}	NI ^{Id}	15±0.5 ^{Fa}
GT	NI ^{Ka}	NI ^{Ia}	NI ^{Ha}	NI ^{Ga}	NI ^{Ha}	NI ^{Fa}	NI ^{Ia}	NI ^{Ha}
P	17±0.57 ^{Ia}	13±0.58 ^{Hb,c}	12±0.59 ^{Fc}	NI ^{Gd}	16±0.51 ^{Ga}	14±0.53 ^{Eb}	12±0.55 ^{Hc}	13±0.55 ^{Gb,c}
R/GT	11±0.55 ^{Jc}	15±0.48 ^{Ga}	12±1 ^{Fb,c}	NI ^{Gd}	NI ^{Hd}	13±0.57 ^{Eb}	NI ^{Id}	16±0.5 ^{E,Fa}
R/P	21±0.58 ^{Ga}	17±0.40 ^{Fb}	15±0.57 ^{Ec}	NI ^{Gd}	20±0.53 ^{Ea}	17±0.15 ^{Db}	15±0.29 ^{Gc}	17±0.5 ^{Eb}
Antibiotic								
CPF	26±0.57 ^{D,Ee}	40±0.54 ^{Aa}	40±0.57 ^{Aa}	31±0.76 ^{Ac}	36±0.57 ^{Bb}	30±1 ^{Ac,d}	28±1 ^{Dd,e}	26±0.58 ^{Ce}
TMF/ SMT	NT ^{Kb}	NI ^{Ib}	NT ^{Hb}	NT ^{Gb}	NT ^{Hb}	NT ^{Fb}	21±0.57 ^{Fa}	NT ^{Hb}
PCL	32±1 ^{Ba}	16±0.48 ^{F,Gc}	NT ^{He}	11±1 ^{Fd}	16±0.57 ^{Gc}	13±0.48 ^{Ed}	29±1 ^{Db}	31±1.05 ^{Ba,b}
CDM	22±0.76 ^{F,Ga}	NI ^{Ic}	NT ^{Hc}	NT ^{Gc}	18±0.53 ^{Fb}	NT ^{Fc}	NT ^{Ic}	NT ^{Hc}
CTX	28±1.05 ^{C,Db}	32±0.46 ^{Ba}	32±0.58 ^{Ba}	22±0.53 ^{C,Dc}	27±0.29 ^{Cb}	21±1.05 ^{Cc}	14±1.05 ^{Gc}	31±0.29 ^{Ba}
IPN	40±1 ^{Ab}	22±1 ^{Dd}	27±0.57 ^{Cc}	26±0.55 ^{Bc}	42±1 ^{Ab}	26±0.58 ^{Bc}	46±0.55 ^{Aa}	42±0.5 ^{Ab}
GTC	24±0.76 ^{E,Fa,b}	20±0.48 ^{Ec,d}	22±1.05 ^{Db,c}	20±1.05 ^{Ec,d}	25±0.58 ^{Da}	21±0.58 ^{Cc,d}	26±0.56 ^{Ea}	19±0.5 ^{Dd}
AMX/ CVT	30±0.56 ^{B,Cb}	26±0.55 ^{Cc}	12±1 ^{Fe}	21±1 ^{D,Ed}	26±0.55 ^{C,Dc}	21±1.05 ^{Cd}	44±0.58 ^{Ba}	42±1.05 ^{Aa}
CFX	28±0.57 ^{C,Db}	26±1 ^{Cc}	22±0.46 ^{Dd,e}	23±0.45 ^{Cd}	20±1.05 ^{Ee}	22±0.48 ^{Cd,e}	40±0.48 ^{Ca}	30±1 ^{Bb}

BACTERIA: SA - *S. aureu*; PA - *P. aeruginosa*; EC - *E. coli*; KP - *K. pneumoniae*; LM - *L. monocytogenes*; BC - *B. cereus*; BL - *B. luteus*; PV - *P. vulgaris*

TINCTURE: R – Reishi; GT - Green tea; P – Propolis

ANTIBIOTICS: CPF – Ciprofloxacin; TMF/SMT - Trimethoprim/sulfamethoxazole; PCL – Penicillin; CDM – Clindamycin; CTX – Ceftriaxone; IPN – Ipinem; GTC – Gentamicin; AMX/CVT - Amoxicillin/clavulanate; CFX - Cefoxitin;

^a column values (different capital letters) and row values (different small letters) are significantly different ($p < 0.05$) by Tukey's multiple range test; NI-no inhibition; NT-not tested. *According to the CLSI protocol microorganisms can be consider resistant if the inhibition zone is lower than 14 mm, intermediate or susceptible-dose dependent for the zone 15-19 mm and susceptible for the zone bigger than 20 mm¹¹

CONCLUSION

The chemical composition and mineral content of pure Reishi tincture and those enriched with propolis or green tea were compared. Among all samples, Reishi/propolis tincture contained a significant amount of polyphenols, with the dominant contribution of flavonoids and phenolic acids. Regarding mineral content, P, K, and Na were the most abundant in all tinctures. Reishi/propolis tincture also exhibited the best antibacterial activity compared to the other extracts. Based on reported chemical compositions and antimicrobial activities, these Reishi-based tinctures have promising prospects as food and health supplements. The high levels of polyphenols, flavonoids, phenolic acids, and essential minerals may contribute to antioxidant, immunomodulatory, and general wellness-promoting effects. Enrichment with propolis or green tea further enhances the bioactive profile, potentially supporting cardiovascular, metabolic, and immune health. Therefore, even if the antibacterial activity is moderate compared to commercial products, the combined health benefits and nutritional value make these tinctures valuable candidates for dietary supplementation. Further studies on bioavailability, dosage, and clinical effects would help to fully realize their potential in human health applications.

SUPPLEMENTARY MATERIAL

Additional data are available electronically at the pages of journal website: <https://www.shd-pub.org.rs/index.php/JSCS/article/view/13428>, or from the corresponding author on request.

Acknowledgements: This work was supported by the Republic of Serbia - Ministry of Science, Technological Development and Innovation (Programs for financing scientific research: 451-03-137/2025-03/200133 and 451-03-136/2025-03/200133 assigned to the University of Niš, Faculty of Technology, Leskovac).

ИЗВОД

ТИНКТУРА РЕИШИЈА (*Ganoderma lucidum*) ОБОГАЂЕНА ПРОПОЛИСОМ ИЛИ ЗЕЛЕНИМ ЧАЈЕМ ИЗ СРБИЈЕ: UHPLC-DAD-ESI-MS КАРАКТЕРИЗАЦИЈА, МИНЕРАЛНИ САСТАВ И АНТИБАКТЕРИЈСКА АКТИВНОСТ

САНДРА С. КОНСТАНТИНОВИЋ¹, ДРАГАН З. ТРОТЕР¹, ЈЕЛЕНА Б. ЗВЕЗДАНОВИЋ¹, САЊА М. ПЕТРОВИЋ¹, САША Р. САВИЋ¹, БОЈАНА Р. ДАНИЛОВИЋ¹ И ЈОВАН ЂИРИЋ²

¹Технолошки факултет, Универзитет у Нишу, Булевар ослобођења 124, 16000 Лесковац, ²Тојличка академија ситроковних ситудија, Одсек за пољопривредно - прехрамбене ситудије, Ђирила и Мејодија 1, 18400 Прокуље.

Циљ овог рада био је процена разлика у хемијском саставу и минералном садржају чисте реиши (*Ganoderma lucidum*) тинктуре са тинктурама обogaђеним прополисом или зеленим чајем. Хемијски састав и садржај минерала су анализирани применом течне хроматографије веома високих перформанси у комбинацији са детектором умрежених фотодиода и електроспреј јонизација-масеном спектрометријом (UHPLC-DAD-ESI-MS) и

оптичком емисионом спектрометријом са индуктивно спрегнутом плазмом (ICP-OES), редом. Међу овим узорцима, реиши/прополис тинктура садржи значајну количину полифенола са доминантним уделом флавоноида и фенолних киселина. У погледу минералног састава, Р, К и Na су највише присутни у свим тинктурама. Реиши/прополис тинктура је показала и најбољу антибактеријску активност, при чему су *P. vulgaris* и *P. aeruginosa* биле најосетљивије бактерије.

(Примљено 19. јуна; ревидирано 21. јула; прихваћено 20. новембра 2025.)

REFERENCES

1. S. Wu, S. Zhang, B. Peng, D. Tan, M. Wu, J. Wei, Y. Wang, H. Luo, *Food Sci. Hum. Wellness* **13** (2024) 568 (<https://doi.org/10.26599/FSHW.2022.9250051>)
2. K. Sułkowska-Ziaja, M. Balik, A. Szczepkowski, M. Trepa, G. Zengin, K. Kała, B. Muszyńska, *Diversity* **15** (2023) 882 (<https://doi.org/10.3390/d15080882>)
3. S. Wachtel-Galor, J. Yuen, J.A. Buswell, I. F. F. Benzie, *Ganoderma lucidum (Lingzhi or Reishi): A Medicinal Mushroom*, in *Herbal Medicine: Biomolecular and Clinical Aspects second edition*, I. F. F. Benzie, S. Wachtel-Galor, Ed(s)., CRC Press/Taylor & Francis, Boca Raton, Florida, USA, 2011, pp. 175–200 (<https://doi.org/10.1201/b10787>)
4. A. K. Balasubramaniam, A. Elangovan, M. A. Rahman, S. Nayak, D. Swain, H. P. Babu, A. Narasimhan, V. Monga, *Fitoterapia* **183** (2025) 106526 (<https://doi.org/10.1016/j.fitote.2025.106526>)
5. N. Bhatti, Y. A. Hajam, S. Mushtaq, L. Kaur, R. Kumar, S. Rai, *Discov. Sustain.* **5** (2024) 185 (<https://doi.org/10.1007/s43621-024-00375-3>)
6. A. Sarapa, A. Peter, A. Buettner, H. M. Loos, *Eur. Food Res. Technol.* **251** (2025) 1331 (<https://doi.org/10.1007/s00217-025-04708-y>)
7. J. Sun, S. Dong, J. Li, H. Zhao, *Food Sci. Hum. Wellness* **11** (2022) 1143 (<https://doi.org/10.1016/j.fshw.2022.04.008>)
8. V. Tadić, I. Arsić, J. Zvezdanović, A. Zugić, D. Cvetković, S. Pavkov, *J. Ethnopharmacol.* **199** (2017) 138 (<https://doi.org/10.1016/j.jep.2017.02.002>)
9. S. V. Dončić, J. B. Zvezdanović, S. M. Petrović, S. R. Savić, I. B. Konstantinović, D. Z. Troter, J. M. Jovanović Mirković, S. S. Konstantinović, *Adv. Technol.* **13** (2024) 5 (<https://doi.org/10.5937/savteh2401005D>)
10. S. Savić, S. Petrović, M. Petronijević, A. Cvetanović, Ž. Petronijević, *Adv. Technol.* **8** (2019) 27 (<https://doi.org/10.5937/SavTeh1901027S>)
11. CLSI Performance Standards for Antimicrobial Susceptibility Testing. 30th ed. CLSI supplement M100, Wayne PA: Clinical and Laboratory Standards Institute, 2020
12. D. H. Ryu, J. Y. Cho, N. B. Sadiq, J.-C. Kim, B. Lee, M. Hamayun, T. S. Lee, H. S. Kim, S. H. Park, C. W. Nho, H.-Y. Kim, *Food Chem.* **335** (2021) 127645 (<https://doi.org/10.1016/j.foodchem.2020.127645>)
13. R. J. Robbins, *J. Agric. Food. Chem.* **51** (2003) 2866 (<https://doi.org/10.1021/jf026182t>)
14. E. de Rijke, P. Out, W. M. A. Niessen, F. Ariese, C. Gooijer, U. A. Th. Brinkman, *J. Chromatogr. A* **1112** (2006) 31 (<https://doi.org/10.1016/j.chroma.2006.01.019>)
15. Z. Yan, B. Xia, M. H. Qiu, D. L. Sheng, H. X. Xu, *Biomed. Chromatogr.* **27** (2013) 1560 (<https://doi.org/10.1002/bmc.2960>)

16. T.-Q. Chen, C. Yang, J.-G. Wu, J.-Z. Wu, *J. Chem. Pharm. Res.* **8** (2016) 730 (<https://www.jocpr.com/articles/total-triterpenoids-from-the-ultrasoniccirculating-extract-powder-of-ganoderma-lucidum-and-its-antioxidant-activity-in-v.pdf>)
17. C. C. Wyrepkowski, D. L. M. Gomes da Costa, A. P. Sinhorin, W. Vilegas, R. A. De Grandis, F. A. Resende, E. A. Varanda, L. C. dos Santos, *Molecules* **19** (2014) 16039 (<https://doi.org/10.3390/molecules191016039>)
18. D. H. Markowicz Bastos, L. A. Saldanha, R. R. Catharino, A. C. H. F. Sawaya, I. B. S. Cunha, P. O. Carvalho, M. N. Eberlin, *Molecules* **12** (2007) 423 (<https://doi.org/10.3390/12030423>)
19. MassBank Record, <http://www.massbank.jp> (accessed 13 May 2025)
20. Z. Benayad, C. Gómez-Cordovés, N. E. Es-Safi, *Int. J. Mol. Sci.* **15** (2014) 20668 (<https://doi.org/10.3390/ijms151120668>)
21. H.-Y. Zhao, J.-H. Sun, M.-X. Fan, L. Fan, L. Zhou, Z. Li, J. Han, B.-R. Wang, D.-A. Guo, *J. Chromatogr. A* **1190** (2008) 157 (<https://doi.org/10.1016/j.chroma.2008.02.109>)
22. X. Jiang, Y. Liu, W. Li, L. Zhao, F. Meng, Y. Wang, H. Tan, H. Yang, C. Wei, X. Wan, L. Gao, T. Xia, *PLoS One* **8** (2013) e62315 (<https://doi.org/10.1371/journal.pone.0062315>)
23. J.-P. Cornard, C. Lapouge, *J. Phys. Chem. A* **110** (2006) 7159 (<https://doi.org/10.1021/jp060147y>)
24. F. Pellati, G. Orlandini, D. Pinetti, S. Benvenuti, *J. Pharm. Biomed. Anal.* **55** (2011) 934 (<https://doi.org/10.1016/j.jpba.2011.03.024>)
25. M. Ye, W.-Z. Yang, K.-D. Liu, X. Qiao, B.-J. Li, J. Cheng, J. Feng, D.-A. Guo, Y.-Y. Zhao, *J. Pharm. Anal.* **2** (2012) 35 (<https://doi.org/10.1016/j.jpha.2011.09.009>)
26. J. M. Fuentes-Alventosa, S. Jaramillo, G. Rodríguez-Gutiérrez, P. Cermeño, J. A. Espejo, A. Jiménez-Araujo, R. Guillén-Bejarano, J. Fernández-Bolaños, R. Rodríguez-Arcos, *J. Agric. Food. Chem.* **56** (2008) 6977 (<https://doi.org/10.1021/jf8009766>)
27. F. Sánchez-Rabaneda, O. Jáuregui, I. Casals, C. Andrés-Lacueva, M. Izquierdo-Pulido, R. M. Lamuela-Raventós, *J. Mass Spectrom.* **38** (2003) 35 (<https://doi.org/10.1002/jms.395>)
28. S. I. Falcão, M. Vilas-Boas, L. M. Estevinho, C. Barros, M. R. M. Domingues, S. M. Cardoso, *Anal. Bioanal. Chem.* **396** (2010) 887 (<https://doi.org/10.1007/s00216-009-3232-8>)
29. J. Sun, F. Liang, Y. Bin, P. Li, C. Duan, *Molecules* **12** (2007) 679 (<https://doi.org/10.3390/12030679>)
30. M. F. Ahmad, A. Alsayegh, F. A. Ahmad, M. S. Akhtar, S. S. Alavudeen, F. Bantun, S. Wahab, A. Ahmed, M. Ali, E. Y. Elbendary, A. Raposo, N. Kambal, M. H. Abdelrahman, *Heliyon* **10** (2024) e25607 (<https://doi.org/10.1016/j.heliyon.2024.e25607>)
31. D. S. Ghallab, E. Shawky, M. M. Mohyeldin, A. M. Metwally, R. S. Ibrahim, *J. Adv. Pharm. Sci.* **2** (2025) 76 (<https://doi.org/10.21608/japs.2025.336494.1032>)
32. M. Radeva-Ilieva, S. Stoeva, N. Hvarhanova, K. D. Georgiev, *Foods* **14** (2025) 745 (<https://doi.org/10.3390/foods14050745>)
33. A. A. H. Ali, *J. Trace Elem. Miner.* **4** (2023) 100076 (<https://doi.org/10.1016/j.jtemin.2023.100076>)
34. M. A. Farag, B. Abib, Z. Qin, X. Ze, S. E. Ali, *Curr. Res. Nutr. Food Sci.* **6** (2023) 100450 (<https://doi.org/10.1016/j.crfs.2023.100450>)

35. M. N. Djide, Sartini, L. Rahman, N. Hasyim, *Int. J. Sci. Technol. Res.* **3** (2014) 15
(<http://www.ijstr.org/final-print/jan2014/Antibacterial-Activity-Of-Various-Extracts-From-The-Fruiting-Bodies-Of-Ganoderma-Lucidum-Growing-At-Samanea-Saman-jacq-Merr-Trunk.pdf>)
36. J. Singh, S. Gupta, S. Malviya, B. Ahrwar, *Int. J. Adv. Res.* **2** (2014) 460
(https://www.journalijar.com/uploads/31_IJAR-3328.pdf)
37. W. C. Reygaert, *Front. Microbiol.* **5** (2014) 434
(<https://doi.org/10.3389/fmicb.2014.00434>)
38. S. Ivančajić, I. Mileusnić, D. Cenić-Milošević, *Arch. Biol. Sci.* **62** (2010) 915
(<https://doi.org/10.2298/ABS1004915I>)
39. P. Ristivojević, I. Dimkić, J. Trifković, T. Berić, I. Vovk, D. Milojković-Opsenica, S. Stanković, *PLoS One* **11** (2016) e0157097
(<https://doi.org/10.1371/journal.pone.0157097>)
40. S. Stepanović, N. Antić, I. Dakić, M. Švabić-Vlahović, *Microbiol. Res.* **158** (2003) 353 (<https://doi.org/10.1078/0944-5013-00215>).