



SUPPLEMENTARY MATERIAL TO
**Reishi (*Ganoderma lucidum*) tincture enriched with propolis or green
tea from Serbia: UHPLC-DAD-ESI-MS characterization, mineral
content and antibacterial activity**

SANDRA S. KONSTANTINOVIĆ^{1*}, DRAGAN Z. TROTTER¹, JELENA B.
ZVEZDANOVIĆ¹, SANJA M. PETROVIĆ¹, SAŠA R. SAVIĆ¹, BOJANA R. DANILOVIĆ¹
AND JOVAN ČIRIĆ²

¹University of Niš, Faculty of Technology, Bulevar oslobođenja 124, 16000 Leskovac, Serbia,
and ²Toplica Academy of Vocational Studies, Department of Agricultural and Food Studies,
Ćirila i Metodija 1, 18400 Prokuplje, Serbia.

* Corresponding author. E-mail: sakisandra12@yahoo.com

TABLE S-I. Parameters of the calibration line for the determined elements

Element	λ / nm	R / 1	ROL / mg·L ⁻¹
Ag	224.641	0.99993	0.00532-12
	328.068	0.99995	0.00166-12
As	189.042	0.99982	0.00255-3.6
	197.262	1	0.00825-2.4
B	182.641	0.99999	0.00737-12
	249.773	0.99999	0.00643-12
Ba	233.527	0.99995	0.000115-12
Bi	190.241	0.99991	0.00353-12
	223.061	0.99996	0.00368-12
Ca	183.801	1	1.6-12
	396.847	1	0.0126-2.41
Cd	214.438	0.99989	0.000111-2.4
Ce	418.660	0.999917	0.00118-2.4
Co	228.616	0.99980	0.000327-12
Cr	283.563	0.99998	0.000435-12
Cu	224.700	0.99999	0.000907-12
	324.754	0.99999	0.000259-12
Fe	259.941	0.99995	0.00199-12
	184.950	0.99992	0.000192-12
Hg	194.227	0.99994	0.000579-12
	253.652	0.99986	0.00202-12
In	325.609	1	0.004-12
K	404.721	0.99996	0.156-60
	323.261	1	0.798-12
Li	670.780	0.99997	0.0000574-1.2
Mg	285.213	0.99996	0.00206-12.1
Mn	257.611	0.99997	0.000105-12
Na	330.237	0.99995	0.798-12
	589.592	1	0.00583-1.2
Ni	231.604	0.99994	0.000474-12
P	214.914	0.99995	0.00-120
Pb	220.353	0.99998	0.00616-12
S	180.731	1	0.0029-24
Si	288.158	0.99998	0.00582-60
	251.612	0.99998	0.00124-60
Tl	190.864	0.99998	0.00172-12
Zn	213.856	0.99997	0.000082-12

TABLE S-II. The compounds detected by UHPLC-DAD-MS/MS in the pure Reishi tincture

Peak No.	t_R / min	λ_{max} / nm	$[M+H]^+$, m/z	MS/MS fragment ion peaks, m/z	Compounds assignment	Reference
1	1.11	280, 350	-	-	-	-
2	1.57	306	-	-	-	-
3	2.28	328	$[M+H]^+$ 301	^a 297.7 (100 %), 255	n.i. phenolic acid	Robbins ¹
4	2.95	280, 450	-	-	-	-
5	3.38	327	$[M+H]^+$ 301	^a 280, 264, 254, 245, 187.9 (100 %), 121	n.i. phenolic acid	Robbins ¹
6	5.81	304	$[M+H]^+$ 240	^a 217 (100 %), 199, 195	n.i. phenolic acid	Robbins ¹
7	7.46	333	175	147 (100 %), 131	n.i. phenolic acid	Robbins ¹
8	8.83	260, 366	373	329 (100 %), 305, 263, 178, 151	n.i. flavonoid	Rijke <i>et al.</i> ²
9	10.10	309	401	-	tent. Lucidone A	Yan <i>et al.</i> ³
			$[M+Na]^+$ 425	^a 407, 226, 161	-	
10	10.70	245	569	-	12-acetoxycanoderic acid F	Chen <i>et al.</i> ⁴ ; Ryu <i>et al.</i> ⁵
11	11.00	237, 315	399	370, 353 (100 %), 337, 325, 309, 159	-	-
			$[M+Na]^+$ 423	377 (100 %), 408, 286, 159	-	-
12	11.55	258	531	-	Ganoderic acid G type A	Chen <i>et al.</i> ⁴
				549.8, 547, 527, 510, 497 (100 %)	Ganoderic acid G or methyl lucidenate	Chen <i>et al.</i> ⁴
13	11.65	-	$[M+Na]^+$ 555	537, 527, 508.66 (100 %), 490.62, 365, 283	Ganoderic acid G or methyl lucidenate	Chen <i>et al.</i> ⁴

^aPositive ionization mode

TABLE S-III. The compounds detected by UHPLC-DAD-MS/MS in the Reishi/green tea tincture

Peak No.			[M-H] ⁻ , <i>m/z</i>	MS/MS fragment ion peaks, <i>m/z</i>	Compounds assignment	Reference
1	0.90	-	343	191 (100 %), 169	3- <i>O</i> -galloyl quinic acid	Wyrepkowski <i>et al.</i> ⁶
2	1.13	287	305	261, 219, 179 (100 %), 175, 159, 137	Galocatechin	Markowicz Bastos <i>et al.</i> ⁷
3	1.24	324	229	183, 161 (100 %), 143	n.i.	
4	1.54	277, 337	305	261, 219, 179 (100 %), 175, 167, 137, 125	Epigallocatechin	Markowicz Bastos <i>et al.</i> ⁷
5	1.77	275	-	-	n.i.	
6	2.23	329	-	-	Phenolic acid derivative	Robbins ¹
7	3.10	274	289	245 (100 %), 231, 203, 175, 161, 137	Catechin/Epicatechin	MassBankRecord ^{8a} , Markowicz Bastos <i>et al.</i> ⁷
8	3.55	312	457	-	Epigallocatechin gallate	Markowicz Bastos <i>et al.</i> ⁷
9	4.69	272, 334	563	473, 443, 383, 353 (100 %), 297	Apigenin 6(or 8)-C-xyloside- 8(or 6)-C-glucoside (vicenin 1)	Benayad <i>et al.</i> ⁹
10	5.12	259, 355	771	-	Apigenin 6-C-glucosyl 8-C-(2"- O-dihydroferuloyl)-glucoside	Benayad <i>et al.</i> ⁹
11	5.28	264, 351	609	301 (100 %), 271, 255, 228	Rutin	Standard
12	5.45	271, 349	463	343, 301 (100 %), 271, 243, 151	Quercetin-glycoside	Zhao <i>et al.</i> ¹⁰
13	5.63	266, 350	593	285 (100 %), 255, 241, 199, 185	Kaempferol 3- <i>O</i> -rutinoside	Jiang <i>et al.</i> ¹¹
14	5.80	268, 312	447	285 (100 %), 255, 227, 137	Kaempferol- <i>O</i> -glycoside	Jiang <i>et al.</i> ¹¹ ; Zhao <i>et al.</i> ¹⁰
15	7.45	333	583	539 (100 %), 469, 395, 243	n.i.	
16	8.85	257, 357	373	329 (100 %), 305, 263, 192, 177, 151	n.i.	
17	10.12	308	401	383, 355	Lucidone A	Yan <i>et al.</i> ³

*PR100687/PR100688

TABLE S-IV. The compounds detected by UHPLC-DAD-MS/MS in the Reishi/propolis tincture

Peak No.	t_R /min	[M-H] ⁺ , m/z	MS/MS fragment ion peaks, m/z	Compounds assignment	Reference
1	1.97	311, 282	137	Hydroxybenzoic acid	MassBankRecord ^{8a}
2	2.71	323, 290	179	Caffeic acid	MassBankRecord ^{8b} ; Cornard and Lapouge ¹²
3	4.71	310	163		MassBankRecord ^{8c} ; Pellati <i>et al.</i> ¹³
4	5.20	323, 290	281	Caffeic acid derivative	-
5	5.47	323, 298	303	Dihydroquercetin	MassBankRecord ^{8d} ; Ye <i>et al.</i> ¹⁴
6	5.60	-	623	Isorhamnetin 3-O-rutinoside	Fuentes-Alventosa <i>et al.</i> ¹⁵
7	5.82	257, 340	447	Quercetin-O-rhamnoside	MassBankRecord ^{8e}
8	6.45	322, 290	431	Genistin	Ye <i>et al.</i> ¹⁴
9	6.92	257, 370	301	Quercetin	Pellati <i>et al.</i> ¹³ ; standard
10	7.13	289, 320	285	Pinobanksin-5-methyl-ether	Pellati <i>et al.</i> ¹³
11	7.22	350, 252	315	Quercetin-3-methyl-ether	Pellati <i>et al.</i> ¹³
12	7.33	280	315	n.i.	
13	7.41	310, 293	349	n.i.	
14	7.62	338, 267	269	Apigenin	Sánchez-Rabaneda <i>et al.</i> ¹⁶ ; standard
15	7.81	293, 330	271	Pinobanksin	Pellati <i>et al.</i> ¹³
16	8.17	267, 350	299	Luteolin-methyl-ether	Pellati <i>et al.</i> ¹³
17	8.70	354, 261	283	Galangin-5-methyl-ether	Pellati <i>et al.</i> ¹³
18	9.04	312	449	n.i.	
19	9.38	258, 358	329	Quercetin-dimethyl-ether	Pellati <i>et al.</i> ¹³
20	9.89	325, 300	247	Caffeic acid isoprenyl ester	Falcão <i>et al.</i> ¹⁷
21	10.05	327, 300	269	Caffeic acid benzyl ester	Pellati <i>et al.</i> ¹³
22	10.16	-	247	Caffeic acid isoprenyl ester – isomer	Falcão <i>et al.</i> ¹⁷
23	10.27	291, 356	255	Pinocembrin	Falcão <i>et al.</i> ¹⁷
24	10.53	296	313	Pinobanksin-3-O-acetate	Falcão <i>et al.</i> ¹⁷
25	10.69	345, 266	283	Methoxy-chrysin	Pellati <i>et al.</i> ¹³
26	11.21	325, 297, 250	295	n.i.	
27	11.45	293, 320	327	Pinobanksin-3-O-propionate	Falcão <i>et al.</i> ¹⁷
28	12.10	292, 330	341	Pinobanksin-3-O-butyrate	Pellati <i>et al.</i> ¹³
29	12.54	293, 330	355	Pinobanksin-3-O-pentanoate	Pellati <i>et al.</i> ¹³

^aPR100187; ^bKO000512; ^cPR100685; ^dBML01276; ^ePR100993

Fig. S-1. a) Ganoderic acid G type A




Fig. S-1. b) Gallocatechin




Fig. S-1. Mass (A) and UV-Vis (B) spectra of selected compounds from chromatograms of pure Reishi tincture (a) and Reishi/green tea tincture (b-d). a) Compound No. 12 (Fig. 1a), assigned as ganoderic acid G type A (Table S-I); b) Compound No. 2 (Fig. 1b), assigned as gallocatechin (Table S-II);

Fig. S-1. c) Apigenin-xyloside-glucoside




Fig. S-1. d) Quercetin-glycoside




Fig. S-1. Continued c) Compound No. 9 (Fig. 1b), assigned as apigenin-xyloside-glucoside (Table S-II); d) Compound No. 12 (Fig. 1b), assigned as quercetin-glycoside (Table S-II).

Fig. S-2. a) Caffeic acid




Fig. S-2. b) Dihydroquercetin




Fig. S-2. Mass (A) and UV-Vis (B) spectra of selected compounds from chromatogram of Reishi/propolis tincture. a) Compound No. 2 (Fig. 1c), assigned as caffeic acid (Table S-III);
b) Compound No. 5 (Fig. 1c), assigned as dihydroquercetin (Table S-III);

Fig. S-2. c) Genistin




Fig. S-2. d) Pinocembrin




Fig. S-2. Continued c) Compound No. 8 (Fig. 1c), assigned as genistin (Table S-III); d) Compound No. 23 (Fig. 1c), assigned as pinocembrin (Table S-III).

REFERENCES

1. R. J. Robbins, *J. Agric. Food. Chem.* **51** (2003) 2866 (<https://doi.org/10.1021/jf026182t>)
2. E. d. Rijke, P. Out, W. M. A. Niessen, F. Ariese, C. Gooijer, U. A. Th. Brinkman, *J. Chromatogr. A* **1112** (2006) 31 (<https://doi.org/10.1016/j.chroma.2006.01.019>)
3. Z. Yan, B. Xia, M. H. Qiu, D. L. Sheng, H. X. Xu, *Biomed. Chromatogr.* **27** (2013) 1560 (<https://doi.org/10.1002/bmc.2960>)
4. T.-Q. Chen, C. Yang, J.-G. Wu, J.-Z. Wu, *J. Chem. Pharm. Res.* **8** (2016) 730 (<https://www.jocpr.com/articles/total-triterpenoids-from-the-ultrasonically-circulating-extract-powder-of-ganoderma-lucidum-and-its-antioxidant-activity-in-v.pdf>)
5. D. H. Ryu, J. Y. Cho, N. B. Sadiq, J.-C. Kim, B. Lee, M. Hamayun, T. S. Lee, H. S. Kim, S. H. Park, C. W. Nho, H.-Y. Kim, *Food Chem.* **335** (2021) 127645 (<https://doi.org/10.1016/j.foodchem.2020.127645>)
6. C. C. Wyrepkowski, D. L. M. Gomes da Costa, A. P. Sinhorin, W. Vilegas, R. A. De Grandis, F. A. Resende, E. A. Varanda, L. C. dos Santos, *Molecules* **19** (2014) 16039 (<https://doi.org/10.3390/molecules191016039>)
7. D. H. Markowicz Bastos, L. A. Saldanha, R. R. Catharino, A. C. H. F. Sawaya, I. B. S. Cunha, P. O. Carvalho, M. N. Eberlin, *Molecules* **12** (2007) 423 (<https://doi.org/10.3390/12030423>)
8. MassBank Record, <http://www.massbank.jp> (accessed 13 March 2020)
9. Z. Benayad, C. Gómez-Cordovés, N. E. Es-Safi, *Int. J. Mol. Sci.* **15** (2014) 20668 (<https://doi.org/10.3390/ijms151120668>)
10. H.-Y. Zhao, J.-H. Sun, M.-X. Fan, L. Fan, L. Zhou, Z. Li, J. Han, B.-R. Wang, D.-A. Guo, *J. Chromatogr. A* **1190** (2008) 157 (<https://doi.org/10.1016/j.chroma.2008.02.109>)
11. X. Jiang, Y. Liu, W. Li, L. Zhao, F. Meng, Y. Wang, H. Tan, H. Yang, C. Wei, X. Wan, L. Gao, T. Xia, *PLoS One* **8** (2013) e62315 (<https://doi.org/10.1371/journal.pone.0062315>)
12. J.-P. Cornard, C. Lapouge, *J. Phys. Chem. A* **110** (2006) 7159 (<https://doi.org/10.1021/jp060147y>)
13. F. Pellati, G. Orlandini, D. Pinetti, S. Benvenuti, *J. Pharm. Biomed. Anal.* **55** (2011) 934 (<https://doi.org/10.1016/j.jpba.2011.03.024>)
14. M. Ye, W.-Z. Yang, K.-D. Liu, X. Qiao, B.-J. Li, J. Cheng, J. Feng, D.-A. Guo, Y.-Y. Zhao, *J. Pharm. Anal.* **2** (2012) 35 (<https://doi.org/10.1016/j.jpba.2011.09.009>)
15. J. M. Fuentes-Alventosa, S. Jaramillo, G. Rodríguez-Gutiérrez, P. Cermeño, J. A. Espejo, A. Jiménez-Araujo, R. Guillén-Bejarano, J. Fernández-Bolaños, R. Rodríguez-Arcos, *J. Agric. Food. Chem.* **56** (2008) 6977 (<https://doi.org/10.1021/jf8009766>)
16. F. Sánchez-Rabaneda, O. Jáuregui, I. Casals, C. Andrés-Lacueva, M. Izquierdo-Pulido, R. M. Lamuela-Raventós, *J. Mass Spectrom.* **38** (2003) 35 (<https://doi.org/10.1002/jms.395>)
17. S. I. Falcão, M. Vilas-Boas, L. M. Estevinho, C. Barros, M. R. M. Domingues, S. M. Cardoso, *Anal. Bioanal. Chem.* **396** (2010) 887 (<https://doi.org/10.1007/s00216-009-3232-8>).