

1 **Embracing green chromatography principles in perindopril,**
2 **amlodipine and indapamide drug mixture analysis using**
3 **β -cyclodextrin modified mobile phase**

4 HUSEINATU OSMAN*, JEVREM STOJANOVIĆ, ANA PROTIĆ, MIRA ZEČEVIĆ
5 and BILJANA OTAŠEVIĆ**

6 *Department of Drug Analysis, University of Belgrade – Faculty of Pharmacy,*
7 *Vojvode Stepe 450, 11221 Belgrade, Serbia*

8 (Received 4 December 2025, revised 6 February, accepted 11 March 2026)

9 *Abstract:* Raising the level of environmental awareness in the field of liquid
10 chromatography is considered indispensable while the use of β -cyclodextrin, as
11 additive, in a mobile phase is promising strategy in this regard. This study pres-
12 ents a method development in line with ICH Q14 regulatory requirements for
13 introducing sustainability and method life-cycle management to separate com-
14 ponents of a cardiovascular multi-drug tablet formulation. At the beginning, the
15 analytical method target profile was defined, separation of perindopril, amlod-
16 ipine, and indapamide in a shortest possible analytical run time. Following risk
17 analysis pointed out that the mobile phase constituents represent the critical
18 method parameters affecting the chromatographic analyses. Design of experiments
19 methodology and desirability function calculation was employed to simultane-
20 ously optimize the levels of concentration of β -cyclodextrin solution, pH value
21 and acetonitrile content in the mobile phase investigated in the ranges 5–15 mM,
22 4.0–6.0 and 20–30 vol. %, respectively. The optimal chromatographic conditions
23 consisted of 10 mM β -cyclodextrin (pH 5.4) and acetonitrile in the volume ratio
24 70:30, 2 mL min⁻¹ flow rate, RP-18e column kept at 25 °C, 215 nm detection
25 wavelength, and 10 μ L injection volume. The eco-friendliness of the method was
26 assessed using the AGREE tool indicating a green and sustainable method was
27 successfully developed.

28 *Keywords:* HPLC method development; design of experiments; desirability func-
29 tion; cardiovascular multi-drug tablet formulation; AGREE assessment tool.

*,** Corresponding authors. E-mail: (*)oshusei@gmail.com;
(**)biljana.otasevic@pharmacy.bg.ac.rs
<https://doi.org/10.2298/JSC251204013O>

30

INTRODUCTION

31

32

33

34

35

36

Achieving sustainability in the domain of pharmaceutical sciences involves integrating environmental and social considerations from the earliest stages of development through to end-of-life management. This includes using green chemistry principles, developing eco-friendly analytical methods, designing sustainable manufacturing processes and optimizing drug product formulations, packaging and distribution to reduce waste and energy consumption.¹

37

38

39

40

41

42

43

44

45

Multi-drug formulations have increasingly become a significant challenge in the field of drug analysis due to the complexity and resource demanding requirements associated with resolving these compounds. This is attributed to the growing interest in rationally designed multi-target drugs, also known as multimodal drugs, network therapeutics or designed multiple ligands. These drugs have emerged as an appealing drug discovery paradigm over the past decade to address diseases with complex etiologies and those exhibiting substantial drug resistance.^{2,3} Multi-drugs are also means to sustainable environment as less resources are used in their manufacture and consecutive quality control analysis.

46

47

48

49

50


51

52

53

54

Multi-component drug formulation used in this research comprised of a mixture of perindopril *tert*-butylamine (also referred as perindopril erbumine), amlodipine besylate and indapamide active pharmaceutical substances which structures are presented in Fig. 1. Their respective drug formulation has been successfully used for the treatment of cardiovascular diseases, considering that perindopril is an angiotensin-converting enzyme inhibitor that lowers blood pressure by reducing sodium and water retention,⁴ amlodipine besylate is a calcium channel blocker that lowers blood pressure by relaxing blood vessels⁵ and indapamide is a diuretic that causes water elimination hence reducing the pressure inside blood vessels.⁶



55

56

57

Fig. 1. Chemical structures of: A) perindopril erbumine, B) indapamide and C) amlodipine besylate.

58 In the realm of drug analysis, high pressure liquid chromatography (HPLC)
59 stands as the de facto gold standard. Among the various HPLC techniques, reversed
60 phase high performance liquid chromatography (RP-HPLC) predominates, com-
61 prising approximately 75 % of the reported methodologies. Historically, the technique
62 has predominantly utilized acetonitrile as preferred organic solvent in the mobile
63 phase. At the same time, contemporary scientific advancements, particularly the
64 green analytical chemistry (GAC) concept, emphasize the primary objective of
65 reducing or eliminating the utilization and production of substances that pose risks
66 to human health or the environment, as delineated by Anastas.¹ It is evident now
67 that HPLC inherently lacks environmental friendliness due to the substantial quan-
68 tities of solvents employed, which ultimately become waste at the end of the ana-
69 lysis. These wastes become more hazardous to the environment as well as the
70 health and wellbeing of analysts involved if the solvents used are toxic. In the light
71 of the growing need to achieve the global Sustainable Development Goals, SDG
72 9.4, which aims to enhance research and upgrade industrial technologies, particu-
73 larly in developing countries, by 2030. SDG 12.4–6 also focuses on the environ-
74 mentally sound management of chemicals and waste, aiming to reduce their release
75 and minimize adverse impacts on human health and the environment.⁷ However,
76 new and promising strategies highlighted as green chromatography principles that
77 rely on the adjustment of the mobile phase composition emerge, thereby enhancing
78 sustainability of the HPLC method without compromising its performance.^{8–11}
79 There have been several assessment tools for measuring such achievement of a
80 method, such as the GAPI, ESA and NEMI index. However, recently released
81 AGREE tool incorporated in an open access software outstands out for his notable
82 and relevant advantages. In addition to the ecological criteria evaluated by other
83 tools, AGREE considers the number of analytes determined in a single run, sample
84 throughput, automation, and the use of chemicals from renewable sources.^{12,13}

85 The inclination towards introducing GAC concept in the field of drug analysis
86 and the literature review of the reported HPLC methods used for the quality control
87 of ternary drug combination containing perindopril erbumine, amlodipine besylate
88 and indapamide revealing that these methods have predominantly utilized exten-
89 sive amounts of toxic organic solvents, indicated that it is necessary to offer analytical
90 procedure improvement by means of development of an alternative sustainable and
91 eco-friendly HPLC method.^{14,15}

92 Cyclodextrins (CDs) are semi-natural compounds derived from starch (a ren-
93 ewable resource) through enzymatic conversion. Their diverse applications in
94 chemistry, pharmaceuticals, food and cosmetics stem from their non-toxicity and
95 cost-effectiveness.¹⁶ When CDs are used as additives into the mobile phases in
96 HPLC analysis they could reduce the organic solvent–water ratio without compro-
97 mising selectivity or resolution. The potential usefulness of CDs in HPLC separ-
98 ations comes out of the ability of CDs to form inclusion complexes with the guest

99 molecules thus affecting their retention. The efficiency of complexation depends
100 on the structural compatibility between the CD and the guest molecule. The height
101 and internal diameter of the CD cavity are determined by the number of glucose
102 units. α -CD has a lower internal diameter compared to β -CD and γ -CD, enabling
103 them to incorporate low molecular weight compounds with aliphatic chains, β -CD
104 can accommodate heterocyclic and aromatic compounds, spanning a wide range
105 of active pharmaceutical substances (APIs), while γ -CD can accommodate com-
106 plex macrocycles and steroids. Among others, β -CDs are the most utilized CDs in
107 pharmaceutical formulations. Furthermore, β -CD's weak adsorption onto C18
108 columns ensures preserving high column performance, facilitates easy washing,
109 and minimizes damage compared to other CDs. At the same time, all CDs possess
110 a unique advantage over other commonly used mobile phase additives by being
111 transparent within the ultraviolet-visible range predominantly used in detectors in
112 HPLC instruments.^{13,17,18}

113 Sustainability in the field of development of analytical methods may be achieved
114 by using various computer technology related products such as advanced data
115 processing software and strategies that enable reduction of experimental work
116 while maintaining high quality of gathered data. In that respect, the application of
117 Design of Experiments (DoE) methodology has been recognized in the field of drug
118 analysis over past decades. DoE is a robust structural approach based on the multi-
119 factorial planning of order and number of experiments to be executed thus reducing
120 the use of resources as well the generation of waste, but as the most important fact,
121 enabling better insight into factor interactions and factor response relations. DoE
122 serves as a technique for optimizing multivariate systems, such as liquid chromato-
123 graphy, where various interdependent mechanisms, including column efficiency,
124 retention factor, ionization efficiency and ion suppression, analytes' solubility, sig-
125 nificantly impact the analysis.¹⁹⁻²¹ At the same time, according to the International
126 council for harmonization of technical requirements for pharmaceuticals for
127 human use (ICH) Q14 guideline, adopting smarter method development approach
128 supported by principles of quality risk management, enhances the reliability of
129 analytical methods.²² Developed methods also undergo the systematic process of
130 method validation in accordance with ICH Q2(R1/R2)²³ guideline which is con-
131 sidered as necessary to ensure that the developed method is fit for its intended use.

132 Having all this in mind, the aim of this study was focused on the DoE
133 supported development and validation of a HPLC method with β -CD-modified
134 mobile phase as an eco-friendly HPLC alternative for the separation of the three
135 APIs, perindopril erbumine, amlodipine besylate and indapamide from a commercially
136 available tablet formulation.

137

EXPERIMENTAL

138

Chemicals and reagents

139

140

141

142

143

β -CD of 98 % purity was purchased from Acros Organics, USA. An HPLC grade acetonitrile, water and ethanol were purchased from J.T. Baker Inc., USA. Perindopril erbumine, amlodipine besylate and indapamide reference substances were of Ph. Eur. quality. Co-Amlessa tablets containing 4 mg of perindopril erbumine, 10 mg of amlodipine besylate and 1.25 mg of indapamide per one tablet (Krka-Farma, Serbia) were purchased from a local drug store.

144

Chromatographic conditions and equipment

145

146

147

148

149

150

151

152

The experiments were performed on a Vanquish Core 3000 HPLC system (Thermo Fisher Scientific) equipped with quaternary pumps, autosampler, thermostated column department, and PDA detector. Chromatographic data was collected using Chromeleon[®] 7.0 and Chrom Quest 4.2 chromatography data system. Merck Chromolith RP-18e column (100 mm×4.6 mm, macropore size 2 μ m, mesopore size 13 nm) was used for separations. β -CD aqueous solutions were prepared in the concentration range 5–15 mM. The organic part of mobile phase consisted of acetonitrile in range 20–30 vol. %. Column temperature was 25 °C, flow rate 2 mL min⁻¹, injection volume 20 μ L and detection wavelength 215 nm.

153

Stock and working solutions preparation

154

155

156

157

158

159

160

161

162

163

Stock solutions were prepared by dissolving the powder reference substances in 50 vol. % ethanol to attain concentrations of 1 mg mL⁻¹ for each API. Stock solutions were further diluted using the mixture of 10 mM β -CD (pH 5.4) and acetonitrile (70:30 volume ratio) as solvent to prepare one working solution containing 20 μ g mL⁻¹ of perindopril erbumine, 50 μ g mL⁻¹ of amlodipine besylate and 6.25 μ g mL⁻¹ of indapamide to be used within method optimization. A series of five working solutions containing raising concentrations of all APIs were prepared from stock solutions by dilution with the aforementioned solvent to be used within method linearity evaluation. Raising concentrations of working solutions were as follows: 10, 15, 20, 25 and 30 μ g mL⁻¹ for perindopril erbumine, 25.0, 37.5, 50.0, 62.5 and 75 μ g mL⁻¹ for amlodipine besylate and 3.125, 4.688, 6.250, 7.813 and 9.325 μ g mL⁻¹ for indapamide.

164

Sample preparation

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

The stock sample solution was prepared with the appropriate amount of powdered tablet mass containing 4 mg of perindopril erbumine, 10 mg of amlodipine besylate and 1.25 mg of indapamide and transferring it to a 50 mL volumetric flask. The volumetric flask was filled with 50 vol. % ethanol and the solution was filtered. Working sample solution was prepared by transferring of 2.5 mL of stock solution in 10 mL volumetric flask and dilution with the aforementioned solvent to attain concentrations 20, 50 and 6.25 μ g mL⁻¹ for perindopril erbumine, amlodipine besylate and indapamide, respectively, denoted as 100 % concentration level with respect to the declared value for each API. This sample solution was used both for method optimization and method repeatability test (6 sample solutions repetitive analysis) and intermediate method precision test (6 working solutions repetitive analysis performed on three separate days by different analysts) within method validation procedure. The similar procedure was applied for preparation of test sample solutions for accuracy test within method validation procedure with the difference that respective volumes of stock solution were transferred in a 50 mL volumetric flask, mixed with placebo and filled to volume with the aforementioned solvent to attain concentrations 80, 200 and 25 μ g mL⁻¹ for perindopril erbumine, amlodipine besylate and indapamide, respectively. This solution was filtered and diluted with the same solvent to attain test solution

181 containing 16, 40 and 5.00 $\mu\text{g mL}^{-1}$ for perindopril erbumine, amlodipine besylate and indap-
182 amide, respectively, representing 80 % concentration level for each API. Similarly, test solu-
183 tions containing 20, 50 and 6.25 $\mu\text{g mL}^{-1}$ representing 100 % concentrations level and 24, 60
184 and 7.50 $\mu\text{g mL}^{-1}$ representing 120 % concentration level for perindopril erbumine, amlodipine
185 besylate and indapamide, respectively, were prepared. The test sample solutions were prepared
186 in triplicate for each of these concentration levels.

187 *Method optimization*

188 The influence of experimental factors on the retention behavior of the analytes was
189 assessed with experimental design whose plan was constructed as well as obtained results pro-
190 cessing using multiple regression analysis was performed using Design Expert 11.0.0 software
191 (Stat-Ease Inc., USA). The Box–Behnken response surface design was used comprising of a
192 total of 15 runs within which 3-level factors are varied in a predefined manner involving factor
193 levels at the midpoints of the edges of the experimental space and level positioned at the center
194 of the experimental space. The center point experiment was repeated 3 times in order to evaluate
195 the experimental error.^{15,16} Simultaneous optimizations of multiple responses or multi-objective
196 method optimization, was also performed using the same software upon setting the specific
197 numerical target values for each of the observed responses. Firstly, individual desirability func-
198 tions are calculated with a value of 1 for the most ideal outcome for each of the observed res-
199 ponses reaching its target value and 0 for the least desirable outcome. Afterwards, the global
200 desirability function was calculated combining the results of individual desirability functions
201 assigning an overall score also expressed in a range of values 0–1. The overall desirability func-
202 tion represents a compromise and points out to the optimal experimental setup for which all
203 observed responses are the closest possible to their individual target value.

204 *Greenness assessment*


205 Using the AGREE assessment scale, the method was graded for its greenness according to
206 a twelve-point metric system adopted from the 12 GAC principles. Each of the 12 input variables
207 is transformed into a scale in the 0–1 range, and the final result is the product of the assessment
208 results for each GAC principle. The output is a clock-like graph, with the overall score and color
209 representation generated using AGREE calculator (University of Vigo, Spain).

210 RESULTS AND DISCUSSION

211 According to the analytical method development strategy described within
212 ICH Q14 recommendations, the first step is to define the target analytical profile
213 of the new method. The optimization goals were accordingly aligned for the devel-
214 oped method to demonstrate satisfactory separation of adjacent peaks achieved
215 within the least possible total analysis time. The observed chromatographic respon-
216 ses recognized as critical method attributes for achieving these optimization goals
217 (CMAs) were retention factor (k) of the last eluting compound from the mixture,
218 indapamide, as well as measure of separation, selectivity factors between all peak
219 pairs (according to the observed elution order, the selectivity factors were: α_{1-2}
220 between amlodipine and perindopril peaks and α_{2-3} between perindopril and inda-
221 pamide peaks).

222 Multiple factors may affect the retention properties of the APIs under review.
223 They may be classified as analysts, samples, mobile phase, column, instrument and

224 detection related factors, as presented in Ishikawa or fishbone diagram (Fig. 2). The
 225 factor classification followed with constant, noise and experimental (CNX) eval-
 226 uation as a risk analysis approach, enables insight into the critical method para-
 227 meters (CMPs) of the analytical method.²¹ The CNX risk-based approach defines
 228 the method parameters that should be kept under control, at constant level (marked
 229 with yellow color), which ones may be disregarded as noise factors (colored in
 230 green) and which parameters require detail experimental evaluation (colored in
 231 red). From the perspective of the GAC concept, special attention must be paid to
 232 materials, energy and waste. Factors that influence solvent consumption and directly
 233 affect the amount of generated wasted include the composition of the mobile phase.
 234 Therefore, the selection of suitable solvents is considered the most important step
 235 in the intended method development process.



236
237

Fig. 2. Classification of important analytical method aspects with risk-assessment.

238 C18 reversed-phase stationary phase and the type of chromatographic column
 239 was selected before the method is optimized, based on previous experience and
 240 recommendations from the literature.¹⁸ The stationary phase of monolithic columns
 241 is composed of highly porous continuous silica network, with macro- and mesopores.
 242 Macropores are 2 μm in size, and they are responsible for low resistance towards
 243 mobile phase flow. Therefore, monolithic columns are compatible with a high
 244 mobile phase flow rate, even up to 10 mL min^{-1} , accompanied by low pressure in
 245 the system. On the other hand, mesopores are smaller in comparison to macropores
 246 and they account for a huge active surface, approximately 300 $\text{m}^2 \text{g}^{-1}$, which enables
 247 efficient chromatographic separation. Allowing faster mobile phase flow rates and
 248 thus shortening the duration of chromatographic analyses, together with its com-
 249 patibility with highly viscous mobile phases, such as CD-modified mobile phases,
 250 made the monolithic column the ideal choice for the separation of the intended
 251 drug mixture. With reference to other studies dealing with the use of β -CD in
 252 HPLC analysis, a study performed by Đajić *et al.*¹⁸ recommended the use of β -CD
 253 aqueous solutions in a concentration range of 5–15 mM. The pH range for intended
 254 β -CD solutions was considered to be aligned with the $\text{p}K_{\text{a}}$ values of the APIs. It

255 was considered reasonable to for perindopril being the least lipophilic compound,
 256 to select a pH range where he is present in an undissociated form (pK_a values of
 257 perindopril, indapamide and amlodipine are 3.79, 8.85 and 9.60, respectively),
 258 while for other two compounds being very lipophilic to select pH value where they
 259 will have smaller affinity towards reversed-phase stationary phase ($\log P$ values of
 260 perindopril, indapamide and amlodipine are 0.63, 2.52 and 2.20, respectively).
 261 Therefore, the pH range of pH 4.0–6.0 was selected. To provide an appropriate
 262 total analytical run time, a relatively small amount of acetonitrile (20–30 vol. %)
 263 was used to complete the mobile phase. The relationship between selected CMPs
 264 whose levels were investigated inside experimental space bordered as described
 265 and CMAs was investigated using DoE supported regression analysis. The plan of
 266 experiments according to Box–Behnken response surface design together with
 267 results obtained for observed responses is presented in Table I.

268 TABLE I. DoE plan of experiments and results obtained for observed responses

β -CD concentration, mM	Acetonitrile content, vol. %	pH	α_{1-2}	α_{2-3}	k
15	20	5	11.388	7.834	13.024
15	25	6	3.301	8.378	6.992
5	20	5	7.097	2.589	5.030
15	25	4	1.945	5.474	4.392
10	25	5	5.087	8.060	7.171
10	25	5	3.884	6.668	6.234
5	30	5	7.920	3.174	4.158
15	30	5	2.507	6.610	4.222
5	25	4	2.812	6.783	7.325
10	25	5	3.693	7.557	7.270
10	30	6	3.191	7.574	4.089
10	20	6	4.960	3.478	4.952
10	30	4	2.469	5.953	4.051
10	20	4	2.863	10.806	14.114
5	25	6	4.885	6.100	3.522

269 Processing the data in Design Expert 11.0.0 software, regression mathematical
 270 models were obtained which enabled interpreting the dependence of the selected
 271 responses on the examined factors in their corresponding ranges. All models were
 272 in the form of a quadratic polynomial equation, while the need for response trans-
 273 formation was indicated by the software in some cases: the response α_{1-2} was used
 274 as is, while the response α_{2-3} was used as inverse square root and the response k
 275 was transformed with the power function, as shown in Table II. To facilitate the
 276 presentation of the polynomial equation, investigated factors are coded as follows:
 277 A stands for β -CD concentration (mM), B for acetonitrile content, vol. % and C for
 278 pH value of the aqueous part of the mobile phase. Only coefficients of regression

279 model whose p -values were lower than 0.05 threshold limit indicating their statis-
 280 tical significance to the observed response were presented. The negative value of
 281 a coefficient in a polynomial equation indicates that the observed response dec-
 282 creases with the increase of the investigated factor levels, while the positive value
 283 of a coefficient indicated the same direction of a change of levels of investigated
 284 factors and values of observed responses. According to the size of the absolute
 285 values of coefficients in polynomial equations, it was evident that acetonitrile con-
 286 tent, vol. %, appeared as the most influential factor for all observed responses,
 287 usually resulting in the observed response decrease, while the influence of other
 288 factors as well as the intensity of two-factor interactions changed from response to
 289 response. Regression models were also evaluated according to obtained values of
 290 coefficients of determination (R^2 , adjusted R^2 and R^2 predicted), and the statistical
 291 significance of lack of fit value (Table III). The closeness of coefficients of
 292 determination to 1 and p -values of lack of fit being greater than 0.5 threshold limit
 293 indicated satisfactory ability of mathematical models to describe the observed
 294 chromatography systems and thus may be used for predicting retention properties
 295 of APIs according to predefined optimization goals.^{19,20}

296 TABLE II. Coefficients of the regression models


Coefficient	α_{1-2}	$1/(\sqrt{\alpha_{2-3}})$	$k^{2.36}$
Intercept	4.220	0.370	0.011
A	-0.450	-0.097	-0.005
B	-1.280	-0.021	0.011
C	0.780	0.065	0.005
AB	-2.430	0.023	0.004
AC	-0.180	-0.140	-0.016
BC	-0.340	-0.051	-0.005
A^2	1.440	0.120	0.007
B^2	1.570	-0.004	0.006
C^2	-2.420	0.021	0.008

297 TABLE III. Statistical profile of the regression models

Model	R^2	Adjusted R^2	Predicted R^2	Lack of fit p -value
α_{1-2}	0.9367	0.8228	0.7602	0.2825
$1/(\sqrt{\alpha_{2-3}})$	0.9872	0.9641	0.8281	0.2701
$k^{2.36}$	0.9958	0.9882	0.9775	0.8938

298 3D-response surfaces were constructed and used as visualization tools for
 299 easier understanding of the APIs' retention behavior (Fig. 3). They show the dep-
 300 endence of each of the observed CMAs from the two selected CMPs while the
 301 remaining CMP is kept on a constant level. The analysis of the 3D response sur-
 302 faces indicated that the increase in the acetonitrile content led to a decrease in

303 retention of all APIs. When observing the influence of acetonitrile on the selectivity factors, the wavy appearance of response surfaces indicated the significant presence of factor interactions or mutual influence of acetonitrile in combination with other CMPs. In addition, the increase of β -CD concentration demonstrated a similar influence on retention pointing out that significant complexation between drug guest molecules and β -CD occurred enabling the mobile phase to retain appropriate elution strength.




310
311
312
313

Fig. 3. Response surfaces describing dependencies of observed responses α_{1-2} , $1/(\sqrt{\alpha_{2-3}})$ and $k^{2.36}$, respectively from factors: acetonitrile content (vol. %) and β -CD concentration (mM), A, pH value and β -CD concentration (mM), B, and pH value and acetonitrile content (vol. %), C.

314 The steep slope of the response surfaces describing the variation of β -CD
315 concentration indicated the most significant influence of this factor on all observed
316 responses, while the pH of the aqueous part of the mobile phase was always of
317 lower importance. Interestingly, although the appearance of the response surfaces
318 for the response $1/(\sqrt{\alpha_{2-3}})$ pointed out to very dramatic shifts within investigated
319 ranges of CMPs, its values revealed relatively good separation of amlodipine and
320 indapamide peaks, while other CMPs needed more detail considerations prior to
321 selection of the optimal conditions. Having in mind that the responses α_{1-2} and $k^{2.36}$
322 demonstrated different trends upon the influence of investigated factors, it was
323 difficult to define from the response surfaces what would the optimal solution be.

324 Therefore, numerical multi-objective optimization was performed using the desirability function (D) calculation.¹¹ It is considered that the desired fulfillment of
325 predefined CMAs is reached if the maximal value of D is equal to 1, indicating
326 that the compromise solution is met for which all the optimization goals are as
327 close as possible to the predefined goals. For the proposed method, analytical target
328 profile of the method was achieved using the following chromatographic conditions:
329 10 mM β -CD (pH 5.4) and acetonitrile (70:30 volume ratio), 25 °C column
330 temperature, 215 nm detection wavelength, 2 mL min⁻¹ flow rate and 10 μ L
331 injection volume. In accordance with ICH Q14 concept,²² measures of quality
332 assurance were further considered in order to prove that CMAs will be reached
333 under these conditions with appropriate probability and therefore model coefficients
334 uncertainty were discussed. Low values of standard errors of model coefficients
335 (varying from model to model in range -0.026–0.090) and low correlation of
336 residuals for all regression models (0.182, 0.093 and 0.091, respectively) were
337 noted thus proving appropriate regression model quality.

338
339 The verification chromatogram was recorded both with mixture of reference
340 substances and sample solution under selected optimal chromatographic conditions
341 with a total analytical run time within three min, as shown in Fig. 4.



342
343
344

Fig. 4. Representative HPLC chromatogram recorded using mobile phase composed of 10 mM β -CD (pH 5.4) and acetonitrile (70:30 volume ratio).

345 System suitability test parameters were further evaluated: the asymmetry factor (As) meeting the acceptance criteria $0.8 < As < 1.5$, the number of theoretical
346 plates (N) greater than 2000 and appropriate system precision expressed as percent
347 relative standard deviation value (RSD) being lower than 1 % (Table IV). The
348 HPLC method was validated to prove its suitability for intended use, the quality
349 control of commercially available tablets. Method linearity and range, precision,
350 repeatability and accuracy were tested following the procedure required by ICH
351 guideline.²³ From the data shown in Table IV, it may be seen that all validation
352 parameters met appropriate acceptance criteria. Method linearity was demonstrated
353 by the value of calculated correlation coefficient being greater than 0.998.
354

355 The *RSD* values indicated good method repeatability and intermediate precision
 356 since they were lower than 2 and 3 % in average, respectively. Average percent
 357 Recovery values calculated for 3 concentration levels were in range 98–102 %
 358 indicating good method accuracy.

359 Table IV. Results of method validation and system suitability tests

Parameter	Amlodipine besylate	Perindopril erbumine	Indapamide
Asymmetry factor	1.33	1.45	1.36
Number of theoretical plates	4150	5907	4856
System precision, <i>RSD</i> / %	0.39	0.72	0.12
Linearity range, $\mu\text{g mL}^{-1}$	25–75	10–30	3.125–9.325
Correlation coefficient	0.9997	0.9991	0.9993
Repeatability, <i>RSD</i> / %	1.35	1.25	1.77
Intermediate precision, <i>RSD</i> / %	1.41	1.30	1.56
Accuracy (Recovery, %)	98.53	99.32	100.01

360 In order to evaluate HPLC method greenness and sustainability profile, the
 361 assignment of penalty points to every step of the analytical procedure was performed
 362 in the accordance with the procedure of calculating AGREE score.^{12,13} In that
 363 respect, acetonitrile present in the mobile phase was labeled as danger and so
 364 appropriate penalty points were taken. In contrast, β -CD was considered safe.
 365 Then, it was noted that none of the chemicals used exceeded the amount of 20 mL
 366 per analysis. Afterwards, energy consumption was considered. HPLC instruments
 367 commonly use more than 1.5 kWh or less than 0.5 kWh of electrical currency per
 368 sample, and therefore penalty points are assigned to this technique. The PDA detector
 369 was used, which takes less than 0.5 kWh of electrical energy per sample. The
 370 generated waste was collected and it is possible to be recycled. HPLC system used
 371 is hermetically closed system. Therefore, penalty points accounting for produced
 372 waste are assigned only on the basis of its amount. The positioning of the analytical
 373 instrument was also taken into consideration, which is an inline analysis. This adds
 374 to greenness points as there's is elimination of manual sampling and reduction of
 375 waste. Considering all mentioned, the final AGREE score obtained was 0.7 (Fig.
 376 5) which is higher compared to the values obtained from similar studies analyzing




Fig. 5. AGREE greenness assessment results.

377 the same three APIs (0.45 and 0.5).^{14,15} This method is therefore declared as far
378 superior in its eco-friendliness.

379 CONCLUSION

380 Understanding that in recent times there is a need to develop analytical proce-
381 dures for the drug analysis which are sustainable and eco-friendly is of outmost
382 importance. The contribution of every stakeholder in the industry is much needed
383 towards the achievement of the sustainable development goals, in this regard the
384 present work is the kind contribution of the authors to develop a new greener and
385 sustainable HPLC method for the analysis of perindopril, amlodipine and indap-
386 amide in a ternary mixture. The method has AGREE score 0.7 indicating the most
387 compliant profile to sustainability and GAC principles compared to previous reports.

388 *Acknowledgement.* The research was funded by the Ministry of Science, Technological
389 Development and Innovation, Republic of Serbia through Grant Agreements with University of
390 Belgrade Faculty of Pharmacy, No. 451-03-33/2026-03/200161 and 451-03-34/2026-03/
391 /200161.

392

ИЗВОД

393 ПРИМЕНА ПРИНЦИПА ЗЕЛЕНЕ ХРОМАТОГРАФИЈЕ У АНАЛИЗИ СМЕШЕ
394 ПЕРИНДОПРИЛА, АМЛОДИПИНА И ИНДАПАМИДА КОРИШЋЕЊЕМ
395 β -ЦИКЛОДЕКСТРИНА КАО МОДИФИКАТОРА МОБИЛНЕ ФАЗЕ

396 ХУСЕИНАТУ ОСМАН, ЈЕВРЕМ СТОЈАНОВИЋ, АНА ПРОТИЋ, МИРА ЗЕЧЕВИЋ И БИЉАНА ОТАШЕВИЋ

397 *Капдегра за анализирање лекова, Универзитет у Београду – Фармацеутички факултет,
398 Војводе Степе 450, 11221 Београд*

399 Подизање нивоа еколошке свести у домену развоја метода течне хроматографије је
400 неопходно, при чему употреба β -циклодекстрина као адитива мобилне фазе представља
401 релативно новију стратегију која обећава. Овај рад приказује развој методе усаглашен са
402 захтевима ИСН Q14 смернице за сагледавање одрживости и управљањем животним цик-
403 лусом методе, са циљем да се обезбеди хроматографска анализа вишеккомпонентне таблет-
404 не формулације која се користи у терапији кардиоваскуларних болести. Најпре је дефини-
405 сисан жељени профил методе, постизање добре раздвојености пикова периндоприла, амло-
406 дипина и индапамида у што краћем времену, а затим је урађена анализа ризика које је
407 указала да компоненте мобилне фазе представљају критичне параметре методе који утичу
408 на ток хроматографске анализе. Методологија дизајна експеримената и израчунавање
409 функције пожељних одговора, искоришћени су за истовремену оптимизацију нивоа кон-
410 центрације раствора β -циклодекстрина, рН вредности и удела ацетонитрила у мобилној
411 фази који су испитивани у опсезима редом 5–15 mM, 4,0–6,0 и 20–30 запр. %. Оптимални
412 хроматографски услови укључивали су: 10 mM раствор β -циклодекстрина (рН 5,4) и
413 ацетонитрил у запреминском односу 70:30 при протоку од 2 mL min⁻¹, RP-18e колону
414 загрејану на 25 °C, таласну дужину детекције од 215 nm и запремину узорка од 10 μ L.
415 Процена еколошке прихватљивости методе помоћу AGREE алата је потврдила да је успе-
416 шно развијена зелена и одржива хроматографска метода.

417

(Примљено 4. децембра 2025, ревидирано 6. фебруара, прихваћено 11. марта 2026)

418

REFERENCES

- 419 1. P. Anastas, *Crit. Rev. Anal. Chem.* **29** (2010) 167 (<https://doi.org/10.1080/10408349891199356>)
- 420 2. W. Löscher, *Front. Pharmacol.* **12** (2021) 730257
- 421 (<https://doi.org/10.3389/fphar.2021.730257>)
- 422 3. J. J. Lu, W. Pan, Y. J. Hu, Y. T. Wang, *PLOS One* **7** (2012) 40262
- 423 (<https://doi.org/10.1371/journal.pone.0040262>)
- 424 4. S. Julius, J. N. Cohn, J. Neutel, M. Weber, P. Turlapaty, Y. Shen, V. Dong, A. Batchelor,
- 425 H. Lagast. *J. Clin. Hypertens.* **25** (2007) 10 ([https://doi.org/10.1111/j.1524-](https://doi.org/10.1111/j.1524-6175.2004.02440.x)
- 426 [6175.2004.02440.x](https://doi.org/10.1111/j.1524-6175.2004.02440.x))
- 427 5. T. Godfraind, *J. Cardiovasc. Pharmacol. Ther.* **19** (2014) 501
- 428 (<https://doi.org/10.1177/1074248414530508>)
- 429 6. D. B. Campbell, *Curr. Med. Res. Opin.* **8** (1983) 9
- 430 (<https://doi.org/10.1185/03007998309109832>)
- 431 7. United Nations General Assembly, *Transforming our world: the 2030 Agenda for*
- 432 *Sustainable Development*, A/RES/70/1 (2015) (<https://docs.un.org/en/A/res/70/1>)
- 433 8. M. Yabré, L. Ferey, I. T. Somé, K. Gaudin, *Molecules* **23** (2018) 1065
- 434 (<https://doi.org/10.3390/molecules23051065>)
- 435 9. S. El Deeb, *Molecules* **29** (2024) 3205 (<https://doi.org/10.3390/molecules29133205>)
- 436 10. J. Stojanović, J. Krmar, B. Otašević, A. Protić, *Arh. Farm.* **73** (2023) 146
- 437 (<https://doi.org/10.5937/arhfarm73-43479>)
- 438 11. N. Đajić, J. Krmar, J. Stojanović, B. Svrkota, B. Otašević, A. Malenović, A. Protić, *Arh.*
- 439 *Farm.* **74** (2024) 235 (<https://doi.org/10.5937/arhfarm74-49572>)
- 440 12. F. Pena-Pereira, W. Wojnowski, M. Tobiszewski, *Anal. Chem.* **92** (2020) 10076
- 441 (<https://doi.org/10.1021/acs.analchem.0c01887>)
- 442 13. L. A. Yahya, C. Vakh, O. Dushna, O. Kalisz, S. Bocian, M. Tobiszewski, *Anal. Chim.*
- 443 *Acta* **1357** (2025) 344052 (<https://doi.org/10.1016/j.aca.2025.344052>)
- 444 14. K. Patel, *Sep. Sci. Plus* **3** (2020) 175 (<https://doi.org/10.1002/sscp.202000010>)
- 445 15. K. N. Prashant, Y. Pragathi, S. Yenduri, *Green Anal. Chem.* **12** (2024) 100182
- 446 (<https://doi.org/10.1016/j.greeac.2024.100182>)
- 447 16. S. Agatonović-Kustrin, M. Lebete, M. E. Brown, D. W. Morton, B. Glass, *J. Serb. Chem.*
- 448 *Soc.* **81** (2016) 1037 (<https://doi.org/10.2298/JSC150915018A>)
- 449 17. V. González-Ruiz, A. G. León, A. I. Olives, M. A. Martín, J. C. Menéndez, *Green*
- 450 *Chemistry* **13** (2011) 115 (<https://doi.org/10.1039/C0GC00456A>)
- 451 18. N. Maljurić, J. Golubović, B. Otašević, M. Zečević, A. Protić, *Anal. Bioanal. Chem.* **410**
- 452 (2018) 2533 (<https://doi.org/10.1007/s00216-018-0911-3>)
- 453 19. U. A. Thorsteinsdóttir, M. Thorsteinsdóttir, *J. Mass Spectrom.* **56** (2021) 4727
- 454 (<https://doi.org/10.1002/jms.4727>)
- 455 20. S. N. Hidayah, M. Mazidatu lakmam, S. M. F. Bukhari, R. N. Yani, *J. Serb. Chem. Soc.*
- 456 **90** (2025) 1203 (<https://doi.org/10.2298/JSC250124062S>)
- 457 21. B. Svrkota, J. Krmar, F. Petronijević, A. Protić, B. Otašević, *J. Sep. Sci.* **48** (2025) 70136
- 458 (<https://doi.org/10.1002/jssc.70136>)
- 459 22. European medicinal agency, *ICH Q14 Guideline on analytical procedure development*
- 460 (2023) ([https://www.ema.europa.eu/en/ich-q14-analytical-procedure-development-](https://www.ema.europa.eu/en/ich-q14-analytical-procedure-development-scientific-guideline)
- 461 [scientific-guideline](https://www.ema.europa.eu/en/ich-q14-analytical-procedure-development-scientific-guideline))
- 462 23. European medicinal agency, *ICH Q2(R2) Guideline on validation of analytical*
- 463 *procedures* (2024) ([https://www.ema.europa.eu/en/ich-q2r2-validation-analytical-](https://www.ema.europa.eu/en/ich-q2r2-validation-analytical-procedures-scientific-guideline)
- 464 [procedures-scientific-guideline](https://www.ema.europa.eu/en/ich-q2r2-validation-analytical-procedures-scientific-guideline)).