

SUPPLEMENTARY MATERIAL TO
**Biosorption of Congo Red and Methylene Blue by pretreated
waste *Streptomyces fradiae* biomass – Equilibrium, kinetic and
thermodynamic studies**

ZDRAVKA Y. VELKOVA¹, GERGANA K. KIROVA^{1*}, MARGARITA S.
STOYTCHEVA² and VELIZAR K. GOCHEV³

¹Department of Chemical Sciences, Faculty of Pharmacy, Medical University of Plovdiv,
Vasil Aprilov 15-A Blvd., Plovdiv 4002, Bulgaria, ²Instituto de Ingeneria, Universidad
Autonoma de baja California, Blvd. B. Juarez y calle de la Normal s/n 21280 Mexicali Baja
California, Mexico and ³Department of Biochemistry and Microbiology, Faculty of Biology,
Paisii Hilendarski University of Plovdiv, Tzar Asen Str. 24,
Plovdiv 400, Bulgaria

J. Serb. Chem. Soc. 83 (1) (2018) 107–120




Fig. S-1. FTIR spectrum of waste pretreated *Streptomyces fradiae* biomass.

*Corresponding author. E-mail: gergana_kiro@mail.bg




Fig. S-2. FTIR spectrum of *S. fradiae* biomass after MB biosorption.




Fig. S-3. FTIR spectrum of *S. fradiae* biomass after CR biosorption.