

SUPPLEMENTARY MATERIAL TO


Removal of textile dyes from water by TiO₂ nanoparticles immobilized on poly(ϵ -caprolactone) beads and foams

DARKA MARKOVIĆ^{1#}, STOJA MILOVANOVIĆ², MARIJA RADOIČIĆ³, ŽELJKO RADOVANOVIC¹, IRENA ZIZOVIC⁴, ZORAN ŠAPONJIĆ³ and MAJA RADETIC^{2**#}

¹Innovation Center of the Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, Belgrade, Serbia, ²Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, Belgrade, Serbia, ³Institute of Nuclear Sciences "Vinča", University of Belgrade, P. O. Box 522, Belgrade, Serbia and ⁴Faculty of Chemistry, Wroclaw University of Science and Technology, Wybrzeże Wyspianskiego 27, 50–370, Wroclaw, Poland

J. Serb. Chem. Soc. 83 (12) (2018) 1379–1389

(a)



(b)




Fig. S-1. The structure of dye C.I. Acid Orange 7 (a) and C.I. Basic Yellow 28 (b).




Fig. S-2. Images of different PCL samples after the first cycle of illumination of dye (AO7 and BY28) solutions.

*Corresponding author. E-mail: maja@tmf.bg.ac.rs




Fig. S-3. Images of the $\text{PCL}_f\text{-TiO}_2$ photocatalyst after repeated illumination cycles of dye (AO7 and BY28) solutions.