

SUPPLEMENTARY MATERIAL TO  
**Complex effect of *Robinia pseudoacacia* L. and *Ailanthus altissima* (Mill.) Swingle growing on asbestos deposits:  
Allelopathy and biogeochemistry**

FILIP J. GRBOVIĆ<sup>1\*</sup>, GORDANA M. GAJIĆ<sup>2</sup>, SNEŽANA R. BRANKOVIĆ<sup>1</sup>, ZORAN B. SIMIĆ<sup>3</sup>, NENAD L. VUKOVIĆ<sup>3</sup>, PAVLE Ž. PAVLOVIĆ<sup>2</sup> and MARINA D. TOPUZOVIĆ<sup>1</sup>

<sup>1</sup>University of Kragujevac, Faculty of Science, Department of Biology and Ecology, Radoja Domanovića 12, 34000 Kragujevac, Serbia, <sup>2</sup>University of Belgrade, Institute for Biological Research “Siniša Stanković”, Department of Ecology, Bulevar despota Stefana 142, 11060 Belgrade, Serbia and <sup>3</sup>University of Kragujevac, Faculty of Science, Department of Chemistry, Radoja Domanovića 12, 34000 Kragujevac, Serbia

J. Serb. Chem. Soc. 85 (1) (2020) 141–153



Fig. S-1. Study site on map of Serbia (A) and satellite image of asbestos deposits of mine “Stragari” (B).

\* Corresponding author. E-mail: filip.grbovic@pmf.kg.ac.rs



Fig. S-2. Asbestos deposits of mine “Stragari” (A) and woody plant species growing on asbestos deposits: *R. pseudoacacia* (B) and *A. altissima* (C).